

Complete Embedded Minimal Surfaces of Finite Total Curvature

David Hoffman*
Department of Mathematics
University of Massachusetts
Amherst, MA 01003

Hermann Karcher†
Math. Institut
Universitat Bonn
D-5300 Bonn, Germany

July 18, 1994

Contents

1	Introduction	2
2	Basic theory and the global Weierstrass representation	4
2.1	Finite total curvature	14
2.2	The example of Chen-Gackstatter	16
2.3	Embeddedness and finite total curvature: necessary conditions	20
2.3.1	Flux	24
2.3.2	Torque	27
2.3.3	The Halfspace Theorem	28
2.4	Summary of the necessary conditions for existence of complete embedded minimal surfaces with finite total curvature	30
3	Examples with restricted topology: existence and rigidity	32
3.1	Complete embedded minimal surfaces of finite total curvature and genus zero: the Lopez-Ros theorem	36
4	Construction of the deformation family with three ends	42
4.1	Hidden conformal symmetries	45
4.2	The birdcage model	47
4.3	Meromorphic functions constructed by conformal mappings	50
4.3.1	The function T and its relationship to u	51

*Supported by research grant DE-FG02-86ER25015 of the Applied Mathematical Science subprogram of the Office of Energy Research, U.S. Department of Energy, and National Science Foundation, Division of Mathematical Sciences research grants DMS-9101903 and DMS-9312087.

†Partially supported by Sonderforschungsbereich SFB256 at Bonn.

4.4	The function z and the equation for the Riemann surface in terms of z and u	52
4.5	The Weierstrass data	55
4.6	The logarithmic growth rates	59
4.7	The period and embeddedness problems for the surfaces $M_{k,x}$	60
4.8	The details of the solution of the period problem, I. Simplification of the integrals.	63
4.9	The details of the solution of the period problem, II. The monotonicity lemma.	66
4.9.1	Proof of the Monotonicity Lemma	68
5	The structure of the space of examples	69
5.1	The space of complete embedded minimal surfaces of finite total curvature .	72
5.2	Some questions and conjectures	73
6	Finite total curvature versus finite topology	76
6.1	Complete, properly-immersed, minimal surfaces with more than one end . .	76
6.2	Complete embedded minimal surfaces of finite topology and more than one end	77
6.3	Complete embedded minimal surfaces of finite topology with one end	78
7	Stability and the index of the Gauss map	80

1 Introduction

We will survey what is known about minimal surfaces $S \subset \mathbb{R}^3$, which are complete, embedded, and have finite total curvature: $\int_S |K| dA < \infty$. The only classically known examples of such surfaces were the plane and the catenoid. The discovery by Costa [14, 15], early in the last decade, of a new example that proved to be embedded sparked a great deal of research in this area. Many new examples have been found, even families of them, as will be described below. The central question has been transformed from whether or not there are any examples except surfaces of rotation to one of understanding the structure of the space of examples.

Up to this point, *every new example* of a complete embedded minimal surface of finite total curvature has been discovered first by using the global version of the Enneper-Riemann-Weierstrass representation, which is essentially due to Osserman [58, 59]. This involves knowledge of the compact Riemann surface structure of the minimal surface, as well as its Gauss map and other geometric-analytic data. One of our goals is to show how this is done in the case that has been most completely analyzed, namely surfaces with genus ≥ 1 and three topological ends. An important quality of this construction is that the Riemann surface and the meromorphic data are constructed simultaneously under the

assumption of symmetry. Moreover, once this is done, parameters must be found in order to have a well-defined finite total curvature surface. This parameter search is typically done by computer using a combination of relatively simple numerical routines and relatively complex graphics tools ([8, 41]). In many cases a full theoretical analysis, as is done here in Section 4 for the three-ended surfaces of Theorem 3.3 has yet to be carried out. Moreover, solution of the period problem does not at all guarantee that the surfaces are embedded. In Section 5 we present examples of Callahan, Hoffman, Karcher, Meeks and Wohlgemuth that lie in one parameter families containing both embedded and immersed surfaces. In fact the period problem and embeddedness are totally independent issues. There are examples of Weierstrass data meeting all necessary conditions (Proposition 2.4) for embeddedness, for which the period problem is not solvable (a genus-one example with two catenoid ends does not exist but Weierstrass data for such a symmetric surface does) and others for which the period problem is solvable for a family of surfaces which are embedded outside of a compact set of \mathbb{R}^3 , but are not embedded.

The book is organized as follows. In Section 2 we present the basic tools of the subject, the most important of which is the Weierstrass-Enneper representation. In Section 2.2 we describe the construction of Chen and Gackstatter [10], which produces a genus-one surface with the symmetries and end-behavior of Enneper's surface. To our knowledge this complete minimal surface of finite total curvature was the first one explicitly constructed by first specifying a geometric property — in this case, end behavior — and then deriving the necessary Weierstrass data. It is not, of course, embedded but its construction, as presented here has most of the features of the construction of the three-ended examples in Section 4. In Section 2.3, the hypothesis of embeddedness is used to derive relationships between the geometry of the surface and its analytic representation. Propositions 2.3, 2.4 and 2.5 gather together all necessary conditions including the relationship between flux, logarithmic growth rates of the ends and residues of the complex differential of the height function.

In Section 3, we present the few global rigidity theorems that are known. (Theorems 3.1 and 3.4, due to Lopez-Ros [51], Schoen [66] and Costa [16].) We present a proof (in Section 3.1) of the Lopez-Ros theorem, which states that a complete minimal surface of genus zero and finite total curvature must be the plane or the catenoid. Our proof follows that of Perez-Ros [60]. We also state the existence result, Theorem 3.3, for the three-ended, complete, minimal surfaces with genus $k - 1$ and k vertical planes of symmetry ([30]). The details of the construction of these surfaces are presented in Section 4. We include here the estimation of the parameters that solve the period problem when $k > 2$. The values of the parameters that close the periods determine the logarithmic growth rates of the ends of these surfaces. For the surface to be embedded, they must lie in a certain range, which they (happily) do. This is done in Sections 4.5–4.9.

In Section 5, we survey other known examples and discuss what little is known about

the structure of the space of complete embedded minimal surfaces of finite total curvature. Section 5.2 presents some conjectures about this space.

Finite total curvature implies finite topology, even without the additional assumption of embeddedness. The converse is not true; the helicoid is simply connected, nonflat and periodic, so its curvature is infinite, while its topology is finite. Up until recently, the helicoid was the only known embedded minimal surface with finite topology and infinite total curvature. In 1992, we discovered, with Fusheng Wei, a complete embedded minimal surface of genus one with one end — asymptotic to the helicoid — that has infinite total curvature [32, 33]. The details of this construction are outside the scope of this work. However, the extent to which finite topology implies finite total curvature is discussed in Section 6.

Section 7 discusses the index of stability of a complete minimal surface. The basic results [22, 26, 25], the equivalence of finite index and finite total curvature, and the fact that the index is completely defined by the Gauss map, are discussed. There is, as yet, no known relationship between embeddedness and properties of the index. This final section is therefore, strictly speaking, misplaced in this book. However the ideas and techniques may, in the long run, prove useful in the study of embeddedness of minimal surfaces.

Acknowledgements.

The authors would like to thank Jim Hoffman and Ed Thayer at the Center for Geometry, Analysis, Numerics and Graphics, at the University of Massachusetts, Amherst for the preparation of the computer graphics illustrations in this article. Thayer, Peter Norman, and Pascal Romon read earlier versions and gave us useful critical comments. Harold Rosenberg helped clarify some important points in Section 2.3. We are grateful to M. Kotani and S. Nayatani for useful comments on Section 7. Robert Bryant and Michael Wolf gave us many useful comments. The authors enjoyed the hospitality of MSRI where the final work on this paper was completed.

2 Basic theory and the global Weierstrass representation

Let $\mathcal{S}, \langle \rangle$ be an oriented surface with a Riemannian metric, and let ∇ denote its Riemannian connection. For a smooth function $f: \mathcal{S} \rightarrow \mathbb{R}$, we will denote its differential by df and its gradient by $\text{grad } f$. They are related by $Vf = df(V) = \langle \text{grad } f, V \rangle, V \in T\mathcal{S}$, which can be thought of as the defining equation for $\text{grad } f$. If w is a one form, its covariant derivative ∇w is defined by the relation

$$(\nabla_U w)V = Uw(V) - w(\nabla_U V). \quad (2.1)$$

The divergence operator is $\text{div} = \text{tr} \nabla$ and the Laplacian of a smooth function f is given by $\Delta f := \text{div} \text{grad } f$. Note that

$$\langle \nabla_U \text{grad } f, V \rangle = U \langle \text{grad } f, V \rangle - (\nabla_U V) f = (\nabla_U df)(V). \quad (2.2)$$

Hence

$$\Delta f = \Sigma(\nabla_{e_i} df)(e_i). \quad (2.3)$$

The operator $R_{\frac{\pi}{2}}$ on vector fields, which rotates each $T_p M$ by $\frac{\pi}{2}$ in the positive direction, is covariantly parallel: i.e. $\nabla \text{Rot}_{\frac{\pi}{2}} = 0$. This means simply that for any vector fields U, V , $\text{Rot}_{\frac{\pi}{2}}(\nabla_U V) = \nabla_U \text{Rot}_{\frac{\pi}{2}} V$, which can be easily checked. This implies that for a vector field V

$$\text{div} V = 0 \Leftrightarrow \nabla \text{Rot}_{\frac{\pi}{2}} V \text{ is symmetric,}$$

$$(\text{i.e. } \langle \nabla_U \text{Rot}_{\frac{\pi}{2}} V, \tilde{U} \rangle = \langle \nabla_{\tilde{U}} \text{Rot}_{\frac{\pi}{2}} V, U \rangle \text{ for all vector fields } U, \tilde{U}).$$

Associated to a vector field W is the dual one form ω defined by $\omega(U) = \langle U, W \rangle$. One can verify by elementary means that

$$d\omega = 0 \Leftrightarrow \nabla W \text{ is symmetric (i.e. } \langle \nabla_U W, \tilde{U} \rangle = \langle \nabla_{\tilde{U}} W, U \rangle \text{ for all vector fields } U, \tilde{U}).$$

$$\Leftrightarrow W \text{ is locally the gradient of some differentiable function.}$$

Combining these two equivalences and applying them to $W = \text{grad } f$, where $f: \mathcal{S} \rightarrow \mathbb{R}$ is some smooth function, we have

$$\Delta f = \text{div}(\text{grad } f) = 0 \Leftrightarrow \text{Rot}_{\frac{\pi}{2}}(\text{grad } f) \text{ is (locally) the gradient of some function } f^*.$$

Smooth functions $f: \mathcal{S} \rightarrow \mathbb{R}$ satisfying $\Delta f = 0$ are called *harmonic functions*. For a harmonic function f , the locally defined functions f^* , where $\text{grad } f^* = \text{Rot}_{\frac{\pi}{2}}(\text{grad } f)$ are called *conjugate* harmonic functions associated to f . Note that $df^* = -df \circ \text{Rot}_{\frac{\pi}{2}}$. Because $\text{grad } f$ and $\text{grad } f^*$ are orthogonal and have the same length, the locally defined mapping given by (f, f^*) is conformal wherever $\text{grad } f \neq 0$. Thus, $f + i f^*$ is a local conformal coordinate in a neighborhood of any point $p \in \mathcal{S}$ where $df_p \neq 0$.

Suppose our surface \mathcal{S} is given as an oriented immersed surface in \mathbb{R}^3 . Let X denote its position vector; we think of X as an immersion $X: \mathcal{S} \rightarrow \mathbb{R}^3$. \mathcal{S} inherits a Riemannian metric $\langle \cdot, \cdot \rangle$ from \mathbb{R}^3 and its Riemannian connection for vector fields can be considered to be the projection of differentiation in \mathbb{R}^3 ; if U, V are vector fields on \mathcal{S} , $\nabla_U V$ is defined by the relation

$$DX(\nabla_U V) = [U(DX(V))]^T, \quad (2.4)$$

where $[\cdot]^T$ denotes projection onto $DX_p(T_p \mathcal{S})$. For $p \in \mathcal{S}$ let $N(p)$ denote the unit normal to $X(\mathcal{S})$ at $X(p)$. The endomorphism $\text{Rot}_{\frac{\pi}{2}}$ on $T_p \mathcal{S}$ is just the pullback by DX of

counter-clockwise rotation around $N(p)$ in $DX[T_p\mathcal{S}]$. We think of N as the Gauss mapping $N:\mathcal{S} \rightarrow S^2$ and denote its differential by DN . By isometric identification of $T_p\mathcal{S}$ with $DX_p(T_p\mathcal{S})$ and $DX_p(T_p\mathcal{S})$ with $T_{N(p)}S^2$ by rigid translation in \mathbb{R}^3 , we may think of DN_p as an endomorphism of $T_p\mathcal{S}$. As such it is called the *shape operator*, and denoted by S . The eigenvalues of S are called the *principal curvatures*, and its eigenvectors are called *principal directions*. The average of the principal curvatures is the *mean curvature* and is denoted by H .

An important formula with strong consequences for minimal surfaces is:

$$\Delta X = -2HN. \quad (2.5)$$

By ΔX , we mean $(\Delta x_1, \Delta x_2, \Delta x_3)$ where (x_1, x_2, x_3) are the Euclidean coordinates in \mathbb{R}^3 . We will verify (2.5) as follows. Writing $DX = (dx_1, dx_2, dx_3)$ and applying (2.1) coordinatewise, we have

$$\begin{aligned} (\nabla_U DX)(V) &= U(DX(V)) - DX(\nabla_U V) \\ &= U(DX(V)) - [U(DX(V))]^T \quad (\text{by 2.4}) \\ &= [U(DX(V))]^N \quad (\text{where } [\cdot]^N \text{ denotes projection onto } DX(T\mathcal{S})^\perp) \\ &= (U(DX(V)) \cdot N)N. \end{aligned}$$

The scalar factor of N in the last expression is $(\pm, \text{depending on your background})$ the Second Fundamental Form. Because $DX(V) \cdot N = 0$, $U(DX(V)) \cdot N = -DX(V) \cdot DN(U) = -\langle S(U), V \rangle$, where S is the shape operator. Thus

$$(\nabla_U DX)(V) = -\langle S(U), V \rangle N.$$

From (2.2) we conclude that $\Delta X = -(tr S)N$, which is (2.5), as desired.

Remark 2.1 *Since the covariant derivative is most often developed on Riemannian manifolds without a given isometric immersion, we indicate how the above definition is used for practical computations. For example, if W is a vector field in \mathbb{R}^3 we can consider its tangential component*

$$W^T := W - \langle W, N \rangle N$$

along the immersed surface $X(S)$. This gives us also a vector field V on the domain surface S such that $DX(V) = W^T$. Strictly speaking it is the covariant derivative of V that was defined, but one often allows abbreviations, e.g. writing W rather than $W \circ X$, in computations like:

$$\begin{aligned} \nabla_U(W^T) &:= DX(\nabla_U V) \\ &= (U(DX(V)))^T \\ &= (U(W - \langle W, N \rangle N))^T \\ &= (UW)^T - \langle W, N \rangle DN(U). \end{aligned}$$

Minimal Surfaces

A minimal surface, \mathcal{S} , is one for which $H \equiv 0$; from (2.5) we can conclude that the coordinate functions of a minimal surface are harmonic. We may conclude from the discussion above that if x is a coordinate function, $x + ix^*$ defines a conformal coordinate chart in a neighborhood of any $p \in \mathcal{S}$ where $dx \neq 0$. Since X is assumed to be an immersion, we conclude that at every $p \in \mathcal{S}$, at least one of the coordinate functions x_i satisfies $dx_i|_p \neq 0$. Thus minimal surfaces inherit naturally, from their Euclidean coordinates, a Riemann surface structure.

Because $trS = H \equiv 0$ on a minimal surface \mathcal{S} , the Gauss mapping $N: \mathcal{S} \rightarrow S^2$ satisfies $\langle DN(U), DN(V) \rangle = \kappa^2 \langle U, V \rangle$ where $\pm\kappa$ are the principal curvatures, and also reverses orientation. We orient $S^2 \subset \mathbb{R}^3$ with the outward-pointing normal (so that *its* Gauss map is the identity and its principal curvatures $+1$) and let σ be stereographic projection from the north pole to the complex plane considered to be the (x_1, x_2) -plane, also positively oriented. With these choices, σ is orientation-reversing and $g := \sigma \circ N: \mathcal{S} \rightarrow \mathbb{C} \cup \{\infty\}$ orientation-preserving and conformal, whenever $DN \neq 0$. Thus g is a *meromorphic function* on \mathcal{S} , considered now to be a Riemann surface.

So far we have a natural Riemann surface structure on the minimal surface and the Gauss map as a meromorphic function. Now we will recognize a minimal surface as the real part of a holomorphic curve. Recall that each coordinate function x gives rise, locally, to a conjugate harmonic function x^* and globally to a holomorphic differential form

$$dx + idx^* = dx - i(dx \circ Rot_{\frac{\pi}{2}}).$$

Associated to our minimal immersion $X = (x_1, x_2, x_3): \mathcal{S} \rightarrow \mathbb{R}^3$ we also have the “conjugate” minimal immersion $X^* = (x_1^*, x_2^*, x_3^*)$ (which is well defined only on some covering \tilde{S} of S). The relation $dX^* = -dX \circ Rot_{\frac{\pi}{2}}$ shows that X and X^* are locally isometric immersions and that $\Psi := X + iX^*$ satisfies $d\Psi \circ Rot_{\frac{\pi}{2}} = i \cdot d\Psi$. In particular Ψ is holomorphic and

$$X = Re\Psi = Re \int d\Psi = Re \int (dX + idX^*).$$

With respect to any local holomorphic parameter $z = u_1 + iu_2$ we have the usual connection with real differentiation:

$$\begin{aligned} d\Psi &= \Psi' dz \\ &= \left(\frac{\partial X}{\partial u_1} + i \frac{\partial X^*}{\partial u_1} \right) dz = \left(\frac{\partial X}{\partial u_1} - i \frac{\partial X}{\partial u_2} \right) dz. \end{aligned}$$

It is convenient to define for a real-valued differentiable function w

$$w_z = \frac{\partial w}{\partial z} := \frac{1}{2} \left(\frac{\partial w}{\partial u_1} - i \frac{\partial w}{\partial u_2} \right),$$

and extend this function linearly to complex-valued functions:

$$(v + iw)_z = v_z + iw_z.$$

If $f = v + iw$ is holomorphic, $f_z = f'$; the desirability of this identity is the reason for the factor of $\frac{1}{2}$. The Cauchy- Riemann equation for f is $v_z = iw_z$, and extending this operation coordinate-wise to $\Psi = X + iX^*$, we have

$$\begin{aligned}\Psi' &= \Psi_z = (X_z + iX_z^*) \\ &= 2X_z = \frac{DX}{\partial u_1} - i \frac{\partial X}{\partial u_2} \\ &= 2iX_z^*.\end{aligned}$$

We find an additional property by computing

$$(\Psi'^2) = 4X_z^2 = \left| \frac{\partial X}{\partial u_1} \right|^2 - \left| \frac{\partial X}{\partial u_2} \right|^2 - 2i \frac{\partial X}{\partial u_1} \cdot \frac{\partial X}{\partial u_2} = 0,$$

because all holomorphic reparametrizations are conformal. The image $\Psi(\tilde{S})$ in \mathbb{R}^3 is therefore called a holomorphic null-curve. Conversely, we see that the real part of a holomorphic null-curve is *conformally* parametrized and, of course, harmonic with respect to the given parametrization. Since conformal changes of the metric do not change the harmonicity of functions, the real part of a holomorphic null-curve is also harmonic with respect to the induced Riemannian metric and hence minimal.

Having established a firm connection with complex analysis, we now tie in some more geometry by rewriting $d\Psi$ in terms of the Gauss map. This leads to a representation formulated by Enneper, Riemann and Weierstrass. Recall that the Gauss map became a meromorphic function after we distinguished a “vertical” direction for the stereographic projection; we take this to be the x_3 -axis. Now we make full use of the fact that our surface is a complex curve: While quotients of 1-forms on a surface are not functions, the quotients $\frac{d\Psi_1}{d\Psi_3}, \frac{d\Psi_2}{d\Psi_3}$ of holomorphic 1-forms on the complex curve are functions, meromorphic functions on the underlying Riemann surface. Moreover, because of $X_z \cdot X_z = 0$, these functions are quadratically related:

$$\left(\frac{d\Psi_1}{d\Psi_3} \right)^2 + \left(\frac{d\Psi_2}{d\Psi_3} \right)^2 = -1.$$

Finally, since

$$\begin{aligned}\frac{\partial X}{\partial u_1} &= \operatorname{Re}(d\Psi(\frac{\partial}{\partial u_1})) \quad \text{and} \\ \frac{\partial X}{\partial u_2} &= \operatorname{Re}(d\Psi(\frac{\partial}{\partial u_2}))\end{aligned}$$

are orthogonal tangent vectors to the minimal surface, we must be able to compute the normal, i.e., the Gauss map g , from $\frac{d\Psi_1}{d\Psi_3}, \frac{d\Psi_2}{d\Psi_3}$. With a small trick, this procedure can be reversed. Put

$$\begin{aligned}f &:= \frac{-(d\Psi_1 + id\Psi_2)}{d\Psi_3} \\ &= \frac{d\Psi_3}{d\Psi_1 - id\Psi_2} \quad (\text{because } (\frac{d\Psi_1}{d\Psi_3})^2 + (\frac{d\Psi_2}{d\Psi_3})^2 = -1).\end{aligned}$$

Then we have

$$d\Psi = \left(\frac{1}{2}\left(\frac{1}{f} - f\right), \frac{i}{2}\left(\frac{1}{f} + f\right), 1\right)d\Psi_3,$$

which now depends only on one meromorphic function f and one holomorphic differential $d\phi_3$. From now on, we will write $d\Psi_3$ as dh . Recall

$$dh := d\Psi_3 = dx_3 - i dx_3 \circ \text{Rot}_{\frac{\pi}{2}}.$$

We know already that f must be closely related to the Gauss map, in fact, we find that inverse stereographic projection of f , namely

$$(2\text{Re}f, 2\text{Im}f, |f|^2 - 1)/(|f|^2 + 1)$$

is orthogonal to the tangent vectors $\text{Re}(\frac{d\Psi}{d\Psi_3}), \text{Im}(\frac{d\Psi}{d\Psi_3})$ since

$$\left\langle \left(\frac{1}{2}\left(\frac{1}{f} - f\right), \frac{i}{2}\left(\frac{1}{f} + f\right), 1\right), (2\text{Re}f, 2\text{Im}f, |f|^2 - 1) \right\rangle = 0 \in \mathbb{R}.$$

This identifies f as the stereographic projection of the Gauss map of the minimal surface. We now state the “Weierstrass representation theorem” with emphasis on its global formulation due to Osserman [59].

Theorem 2.1 *Suppose S is a minimal surface in \mathbb{R}^3 , M its Riemann surface, $g = \sigma \circ N$ the stereographic projection of its Gauss map. Then S may be represented (up to a translation) by the conformal immersion*

$$X(p) = \text{Re} \int \Phi, \text{ where} \tag{2.6}$$

$$\Phi = (\phi_1, \phi_2, \phi_3) = \left((g^{-1} - g) \frac{dh}{2}, i(g^{-1} + g) \frac{dh}{2}, dh \right). \tag{2.7}$$

Conversely, let M be a Riemann surface, $g: M \rightarrow \mathbb{C} \cup \{\infty\}$ a meromorphic function and dh a holomorphic one-form on M . Then (2.6) and (2.7) define a conformal minimal mapping of some covering of M into \mathbb{R}^3 , which is regular provided the poles and zeros of g coincide with the zeros of dh . The mapping X is well-defined on M if and only if no component of Φ in (2.7) has a real period. That is

$$\text{Period}_\alpha(\Phi) =: \text{Re} \oint_\alpha \Phi = 0 \tag{2.8}$$

for all closed curves α on M .

We now compute the basic geometric quantities of M in terms of g and dh . The metric induced on M by X can be expressed as

$$ds^2 = \frac{1}{2}|\Phi|^2 = \frac{1}{4}(|g| + |g|^{-1})^2 |dh|^2, \tag{2.9}$$

The metric will be complete provided $\int_{\delta} ds = \infty$ for every divergent curve δ on M . The Gauss curvature of this metric may be computed as the ratio of the area stretching of dN to that of dX . Then

$$K = \frac{-4|dg|^2}{(|g| + |g|^{-1})^2 |dh|^2} \cdot \frac{4}{(1 + |g|^2)^2}.$$

The last term is the conformal stretching of stereographic projection. The minus sign comes from the fact that stereographic projection reverses orientation. We may rewrite this as

$$K = \frac{-16}{(|g| + |g|^{-1})^4} \left| \frac{dg/g}{dh} \right|^2. \quad (2.10)$$

To express the shape operator $S = DX \cdot DN$ in terms of g and dh , we compute $X_{zz}dz^2$ and take the real part of the dot product with N :

$$X_{zz}dz^2 = \frac{1}{4}(X_{11} - X_{22} - 2iX_{12}) \cdot (du_1^2 - du_2^2 + 2idu_1du_2).$$

Since in general $X_{ij} \cdot N = -X_i \cdot N_j = -\langle S(\frac{\partial}{\partial u_i}), \frac{\partial}{\partial u_j} \rangle = -s_{ij}$ and $X_{11} \cdot N = -X_{22} \cdot N$ because we are using conformal coordinates on a minimal surface

$$-2\operatorname{Re}((X_{zz} \cdot N)dz^2) = (s_{11}du_1^2 + s_{22}du_2^2 + 2s_{12}du_1du_2). \quad (2.11)$$

The right-hand-side is the quadratic form associated with the shape operator S , expressed in terms of the conformal coordinates (u_1, u_2) , where $z = u_1 + iu_2$. The left-hand-side of (2.11) can be written, using (2.7), as $-\operatorname{Re}(\Phi_z dz \cdot N)$. Using (2.7)

$$\Phi_z dz = \frac{-1}{2}((g^{-1} + g), i(g^{-1} - g), 0) \frac{dg}{g} dh,$$

and the fact that

$$N = \sigma^{-1} \circ g = \frac{2}{1 + |g|^2} (\operatorname{Re} g, \operatorname{Im} g, \frac{|g|^2 - 1}{2}),$$

gives in (2.11) that

$$\Sigma s_{ij} du_i du_j = \operatorname{Re} \left\{ \frac{dg}{g} dh \right\}.$$

For a tangent vector $V = v_1 \frac{\partial}{\partial u_1} + v_2 \frac{\partial}{\partial u_2}$, we may express the Second Fundamental Form (2.11) as the real part of a holomorphic quadratic differential:

$$\langle S(V), V \rangle = \operatorname{Re} \left\{ \frac{dg}{g} (V) \cdot dh(V) \right\}. \quad (2.12)$$

This formula allows us to conclude that a curve c on $X(M)$ is

$$\begin{aligned} \text{asymptotic} &\iff \frac{dg}{g}(\dot{c}) \cdot dh(\dot{c}) \in i \quad . \\ \text{principal} &\iff \frac{dg}{g}(\dot{c}) \cdot dh(\dot{c}) \in \mathbb{R} \quad . \end{aligned} \quad (2.13)$$

For a minimal surface X as in (2.7) and (2.8) with the Weierstrass data $\{g, dh\}$ on M , we make the following definition.

Definition 2.1 *The associate surfaces to X are the minimal immersions X_θ given by the Weierstrass data $\{g, e^{i\theta}dh\}$, $0 \leq \theta \leq \frac{\pi}{2}$, using (2.7) and (2.8). The conjugate surface X^* to X is the associate surface $X_{\frac{\pi}{2}}$ with data $\{g, idh\}$.*

Note that the Weierstrass data $\{e^{i\theta}g, dh\}$ produces the same minimal surface as $\{g, dh\}$ rotated by an angle θ around the vertical axis. One sees immediately that X_θ may not be well-defined on M because the period condition (2.8) is not necessarily satisfied. We have

$$X^* = -Im \int \Phi, \text{ and}$$

$$X_\theta = (\cos \theta)X - (\sin \theta)X^*$$

locally. Clearly X_θ is well-defined on M , for all θ , $0 < \theta \leq \frac{\pi}{2}$, if and only if X^* is well-defined on M . Also $X^{**} = -X$. By definition, all members of the associate family share the same Gauss map and from (2.8) it follows that they are isometric.

The Schwarz Reflection Principle for Minimal Surfaces

A curve on any surface M in \mathbb{R}^3 is a straight line if and only if its geodesic curvature and its normal curvature vanish. A principal curve on M that is not a straight line has the property that it is a geodesic if and only if it lies in a plane orthogonal to the surface. Thus (2.13) can be used on geodesics to identify lines and planar principal curves. From (2.13) we see that principal geodesics of X correspond to straight lines of X^* , and vice versa. The Gauss map along a principal geodesic has values parallel to the plane of this curve, and the same values along the conjugate straight line – so that the line is orthogonal to the plane. The Schwarz reflection principle for minimal surfaces states that minimal surfaces with such curves must possess Euclidean symmetries.

If a minimal surface contains a line segment L , then it is symmetric under rotation by π about L . (If a minimal surface is bounded by a line segment L , it may be extended by rotation by π about L to a smooth minimal surface containing L in its interior.)

If a minimal surface contains a planar symmetry curve, then it is symmetric under reflection in the plane of that curve. (If a minimal surface meets a plane orthogonally on its boundary the surface may be extended by reflection through the plane to a smooth minimal surface with this curve in its interior.)

We will prove this below. Finding a geodesic on a minimal surface satisfying one of the conditions of (2.13) is equivalent to finding a straight line or a planar geodesic, which with the Reflection Principle implies the existence of a Euclidean symmetry.

Suppose a minimal surface S contains a line segment L ; without loss of generality we may assume that L is a portion of the x_3 -axis containing $\vec{0}$. The surface S^* conjugate to S contains a planar geodesic in a horizontal plane. Because the coordinate functions of S are harmonic $z := x_3 + ix_3^*$ defines a conformal parameter in a neighborhood of L . To avoid

confusion, we write $z = u_1 + iu_2$ and we may translate so that $(x_1, x_2, x_3)(0) = \vec{0}$. Then

$$x_3(u_1) = u_1, \quad x_3^*(u_1) = 0$$

$$x_1(u_1) = x_2(u_1) = 0.$$

Thus $x_3 + ix_3^*$ maps a line segment on the real axis to the real axis, while $x_1 + ix_1^*$ and $x_2 + ix_2^*$ map this same segment to the imaginary axis.

The Schwarz Reflection Principle for complex analytic functions states if an analytic function f , defined in a neighborhood of a segment of the real axis, maps the segment of the real axis into a line $\ell \subset \mathbb{C}$ then $f(\bar{z}) = \rho \circ f(z)$, where ρ is reflection in ℓ . (Moreover if f is defined as a one-sided neighborhood of the real segment, the extension of f by $f(\bar{z}) := \rho \circ f(z)$ defines an extension of f analytic on a neighborhood of the real line segment).

Since reflection in the real (resp. imaginary) axis is $z \rightarrow \bar{z}$ (resp. $z \rightarrow -\bar{z}$), we may conclude that

$$x_3 + ix_3^*(\bar{z}) = \overline{x_3 + ix_3^*} \quad \text{by definition}$$

$$x_k + ix_k^*(\bar{z}) = \overline{-x_k + ix_k^*} \quad k = 1, 2.$$

Hence

$$X(\bar{z}) := (x_1, x_2, x_3)(\bar{z}) = (-x_1, -x_2, x_3)(z)$$

$$X^*(\bar{z}) := (x_1^*, x_2^*, x_3^*)(\bar{z}) = (x_1, x_2, -x_3)(z).$$

That is, S is symmetric under rotation about the x_3 -axis, and S^* is symmetric under reflection in the (x_1, x_2) -plane, as claimed. (The statements in parentheses of the Schwarz Reflection Principle for minimal surfaces follow from the statements in parentheses for the classical reflection principle.) This completes the proof of the Schwarz Reflection Principle.

The Catenoid and the Helicoid As a simple illustration, consider the catenoid, which can be represented as $M = \mathbb{C} - \{0\}$ with Weierstrass data $\{g, dh\} = \{z, \frac{dz}{z}\}$. See Figure 2.0.

The mapping $\rho_\theta(z) \rightarrow e^{i2\theta}\bar{z}$ is a reflection in the line L_θ given by $c(t) = te^{i\theta}$. Using (2.9) we see that ρ_θ is an isometry, and therefore its fixed point set $X(L_\theta)$ is a geodesic. The mapping (2.6) is well defined because there is only one homology cycle on which to check (2.8), namely that generated by the cycle $|z| = 1$, and:

$$\begin{aligned} \text{Period}_{|z|=1}(\Phi) &= \text{Re} \int_{|z|=1} \left(\left(\frac{dz}{z^2} - dz \right), i \left(\frac{dz}{z^2} + dz \right), \frac{dz}{z} \right) \\ &= \text{Re}(0, 0, 2\pi i) = \vec{0}. \end{aligned}$$

Along rays $c(t) = te^{i\theta}, t > 0$, $c(t) = e^{i\theta}$ and

$$\frac{dg}{g}(e^{i\theta}) \cdot dh(e^{i\theta}) = \left(\frac{e^{i\theta}}{te^{i\theta}} \right)^2 = t^{-2} \in \mathbb{R}.$$

Figure 2.0 The catenoid and helicoid.

The catenoid is the image of $C - \{0\}$ under the Weierstrass mapping with $g = z$ and $dh = cdz/z$, $c = 1$. The image of $\gamma := \{|z| = 1\}$ is the waist-circle of the catenoid (left). When $c = i$, the Weierstrass mapping is multivalued; its third coordinate has a vertical period of length 2π on γ (center). The multivalued mapping produces the helicoid, which can also be produced by reflection across the boundary lines of the image. For $c = e^{i\theta}$, $\theta \neq 0, \text{ mod } \pi$, the image of γ is a helix and the full image surface is not embedded. For any c , $|c| = 1$, the full image surface is complete and has infinite total curvature unless $c = 1$, in which case we have the catenoid with total curvature equal to -4π .

Hence each ray is a planar geodesic in a vertical plane containing the vector $(\cos \theta, \sin \theta, 0)$. Any two reflections must generate a rotation about a common axis and we can conclude, as we know, that the catenoid is a surface of rotation. Since

$$x_3(z) = \operatorname{Re} \int_{z_0}^z \frac{dz}{z} = \ln \frac{|z|}{|z_0|},$$

we see that the surface diverges to ∞ (resp. $-\infty$) in the vertical direction as $|z| \rightarrow \infty$ (resp. 0).

If we consider the conjugate surface X^* of the catenoid, we have Weierstrass data $\{z, idz/z\}$ on $\mathbb{C} - \{0\}$. But now

$$\operatorname{Period}_{|z|=1} \{\Phi^*\} = \operatorname{Re} \int_{|z|=1} i\Phi = \operatorname{Re}(0, 0, i \ln z) = (0, 0, 2\pi).$$

Hence X^* is *not* well-defined on $\mathbb{C} - \{0\}$. It gives rise to a multi-valued mapping, the image of which we know to be fibred by horizontal lines and invariant under a vertical translation. This surface is, of course, the helicoid. Note that since the helicoid and catenoid are locally conjugate, they are locally isometric. But they are not congruent because the catenoid contains no lines. The Gauss map of the catenoid covers S^2 exactly once, omitting two points. Hence its total curvature is -4π . The helicoid is nonflat and periodic, hence it has infinite total curvature.

In the case of finite total curvature, $\int_M K dA > -\infty$, much more is known about the Weierstrass representation.

2.1 Finite total curvature

The representation Theorem 2.1 can be significantly sharpened when the minimal surface in question is complete and has finite total curvature.

Theorem 2.2 (Osseman, [48, 59]) *Let $X: M \rightarrow \mathbb{R}^3$ be a complete conformal minimal immersion with finite total curvature (not necessarily embedded). Then:*

- i) M is conformally diffeomorphic to $\overline{M}_k - \{p_1, \dots, p_r\}$ where \overline{M}_k is a closed Riemann surface of genus k and p_1, \dots, p_r are points in \overline{M}_k , $r \geq 1$;*
- ii) X is proper;*
- iii) The Gauss map $N: M \rightarrow S^2$, which is meromorphic on M , extends to a meromorphic function on \overline{M}_k ; the holomorphic one-form dh extends to a meromorphic one-form on \overline{M}_k .*

iv) The total curvature is an integer multiple of 4π and satisfies

$$\int_M K da \leq -4\pi(k + r - 1) = 2\pi(\chi(M) - r), \quad (2.14)$$

where k and r are integers defined in Statement i), and $\chi(M)$ is the Euler characteristic of M .

Theorem 2.2 is a prescription for constructing finite total curvature examples. We will give an example of how to use it in Section 2.2.

In what follows, we will assume that $X: M \rightarrow \mathbb{R}^3$ is a complete, regular, conformal minimal immersion that has finite total curvature, and that $M = \overline{M}_k - \{p_1, \dots, p_r\}$, where \overline{M}_k is a compact, connected Riemann surface of genus k .

Definition 2.2 Let D_j be a punctured neighborhood of $p_j \in \overline{M}_k$. We will refer to $X(D_j) = E_j$ as an end of M , and define, for fixed $R > 0$,

$$S_{R,j} = \{q \in S^2 \mid Rq \in E_j\} \quad (2.15)$$

where S^2 denotes the unit sphere in \mathbb{R}^3 .

Theorem 2.3 (Gackstatter [24] Jorge-Meeks [42]) Let $S_{R,j}$ be as defined in (2.15).

i) For each $j = 1, \dots, r$, $S_{R,j}$ converges smoothly as $R \rightarrow \infty$ to a great circle, covered an integral number of times;

ii) Let d_j be the multiplicity of the great circle $\lim_{R \rightarrow \infty} S_{R,j}$. Then

$$\begin{aligned} \int_M K dA &= 2\pi \left(2(1 - k) - r - \sum_{j=1}^r d_j \right) \\ &= 2\pi \left(\chi(M) - \sum_{j=1}^r d_j \right). \end{aligned} \quad (2.16)$$

Note that (2.16) gives an interpretation of the difference between the right- and left-hand sides of (2.14).

Definition 2.3 The total spinning of S is the integer quantity

$$n(S) = \sum_{j=1}^r d_j.$$

It is immediate from the definitions of $n(M)$ and d_j that $n(M) \geq r$, with equality if and only if each end has multiplicity one. This may be interpreted in terms of the end at p_j . We will say that M has an “embedded end at p_j ” provided X embeds some punctured neighborhood of p_j . The end at p_j is embedded if and only if $d_j = 1$, as follows from the definition of d_j in Theorem 2.3 ii).

Remark 2.2 For later use in Section 7 we will need to estimate the degree d , of the Gauss map of a complete minimal surface M in terms of the genus. We have from (2.16)

$$d = (k + \frac{r}{2} - 1 + \frac{n(M)}{2}).$$

If $r \geq 2$ then clearly $d \geq k + 1$. If $r = 1$, then $n(M)$ is just d_1 and must be odd, and $d_1 = 1$, if and only if the end of the surface is embedded. Since $\frac{r}{2} + \frac{n(M)}{2} \geq 1$ with equality if and only if $r = 1$ and $n(M) = d_1 = 1$, it follows that $d \geq k + 1$ on a complete minimal surface, unless it has precisely one end, and that end is embedded. However such an exceptional minimal surface must be a plane, as we will show below. Hence on a complete nonplanar minimal surface

$$d \geq k + 1.$$

Several arguments can be used to show that a complete immersed minimal surface M of finite total curvature with a single embedded end is a plane. The simplest uses the halfspace theorem (Theorem 2.4 below in section 2.3.3), which says that a complete properly immersed minimal surface that lies in a halfspace is a flat plane. Because we have finite total curvature and an embedded end, we can use Proposition 2.3 iv) and Remark 2.4 i) to conclude the same thing as follows. The end in question is asymptotic to the end of a plane or the end of a catenoid. In either case, the surface M lies in a halfspace. If the end is catenoidal it is clear that the plane defining the smallest halfspace containing M is tangent to M , violating the maximum principle. If the end of M is planar then M lies in some slab. But again the boundary planes of the smallest slab are tangent to M , a contradiction, unless the slab has zero thickness. But this means M is a plane.

2.2 The example of Chen-Gackstatter

Chen and Gackstatter [10, 11] were the first to construct an example by explicitly solving the period problem. In retrospect, and from our point of view, what they did was put a handle in Enneper's surface, making it a genus-one surface with one end, while preserving all of the symmetries and end-behavior. See Figure 2. This was the first known complete minimal torus with finite total curvature and one end. We will present this construction, following Karcher [43], Barbosa and Colares [2] and Thayer [69, 70].

Enneper's surface is given by the data

$$g = z \quad dh = z dz \tag{2.17}$$

on \mathbb{C} , the sphere minus the point at infinity. There are no nontrivial closed cycles, so there is no period problem. Moreover, the metric induced by using (2.17) in (2.7) and (2.8) is given as in (2.9) by $ds = \frac{1}{2}(|z|^2 + 1)dz$ and is invariant under the maps $z \rightarrow e^{i2\theta}\bar{z}$, for any fixed θ . The line L_θ through $\vec{0}$ given by $e^{i\theta}t, \infty < t < \infty$, is the fixed-point set of this mapping and so is mapped to a geodesic on Enneper's surface. The normal curvature

Figure 2.2 Enneper's surface (left) and the Chen-Gackstatter Surface (right).

(multiplied by the length squared of the tangent vector) of this geodesic is given, according to (2.12) by $Re\{e^{2i\theta}\}$. By (2.13), L_θ is mapped to a straight line on Enneper's surface if $\theta = \pm\frac{\pi}{4}$, and to a planar geodesic if and only if $\theta = 0, \frac{\pi}{2}$. Assuming we start the integration in (2.6) at 0, this implies that Enneper's surface has the (x_1, x_3) - and the (x_2, x_3) -plane as reflective planes of symmetry, and contains the lines $x_1 \pm x_2 = x_3 = 0$.

The planar symmetry lines cut the surface into four congruent pieces. The symmetry lines themselves cross at $\vec{0}$ and the end point at infinity. In addition, the two straight lines cross at $\vec{0}$ and at the end, and the surface is invariant under rotation about these lines.

We will construct a genus-one surface S with the same symmetries and end-behavior as Enneper's surface. From (2.16) we may conclude that the winding d_1 at the end of Enneper's surface is equal to 3. The two reflections generate a rotation R of order-two about the vertical axis. By (2.16), the total curvature of this surface must be -8π , which is equivalent to saying that the degree of g is two. By the symmetry assumptions, the Gauss map is vertical at the end. By symmetry again, and the fact that the degree of g is two, all the vertical normals are fixed points of both reflections, which means they lie on the vertical axis. We place the intersection of the two lines at $\vec{0}$. Because rotation about the lines on the surface reverses orientation, any axis point, p , of the surface (where the normal must be vertical) other than $\vec{0}$, is paired with $-p$, where g has the same value. There must be at least one such point p on the positive x_3 -axis and we orient the surface so that $g(p) = \infty$. Since $g(\vec{0})$ is vertical, it follows that $g(0) = 0$ and $g = 0$ at the end. There can be no other points of the surface on the vertical axis. We have

$$g(e) = 0, g(\vec{0}) = 0, g(p) = g(-p) = \infty, p = (0, 0, \alpha),$$

where the end is at e on the torus \overline{S} .

Choose a triangulation of \overline{S} whose vertices include $0, \pm p$, and e . This triangulation projects to a triangulation of \overline{S}/R with half the number of faces and half the number of edges. But the number of vertices is $\frac{V}{2} + 2$, where V is the number of vertices in the triangulation of S . Since S is a torus

$$\chi(\overline{S}/R) = \frac{\chi(\overline{S})}{2} + 2 = 2;$$

so S/R is a sphere.

Let $z: S \rightarrow S/R = S^2$ denote projection onto the quotient. The meromorphic function z has degree two and is branched at the points $\{\vec{0}, \pm p, e\} \subset \overline{S}$. Without loss of generality we may assume that $z(\vec{0}) = 0$, $z(p) = 1$ and $z(e) = \infty$. Reflection K in either one of the vertical planes of symmetry induces the same order-two automorphism of S^2 . This automorphism fixes $0, 1, \infty$ and $z(-p)$. We may conclude that this automorphism is complex conjugation $z \rightarrow \bar{z}$ and therefore $z(-p)$ is real. Rotation about either line in S induces an orientation reversing automorphism of S^2 . This automorphism fixes 0 and ∞ while interchanging $\pm p$. Thus rotation induces $z \rightarrow -\bar{z}$ on S^2 . Moreover, $z(-p) = -1$ and the two lines on S are projected by z onto the imaginary axis.

We now have the torus \overline{S} presented as a double covering of S^2 , branched at $0, \infty$ and ± 1 . Therefore, the surface \overline{S} is the Riemann surface

$$w^2 = z(z-1)(z+1).$$

This is the square torus on which z is the Weierstrass \mathcal{P} -function, normalized to have a double pole at $e = (\infty, \infty)$, a double zero at $\vec{0} = (0, 0)$, (and branch points at $\pm p = (\pm 1, 0)$). Introducing $\gamma = z/w$, we have

$$\gamma^2 = \frac{z}{(z-1)(z+1)}. \quad (2.18)$$

The function γ has degree equal to two, and has simple zeros at $\vec{0}$ and e , and simple poles at $\pm p$. Therefore $g = \rho\gamma$ for some nonzero complex constant ρ . The Gauss map is either purely real or purely imaginary along curves of S projecting onto the real axis in the z -plane. This is because these curves lie in the symmetry planes $x_2 = 0$ and $x_1 = 0$. Because γ is real on the axis it follows that ρ is either real or imaginary. After a rotation of $\pi/3$, if necessary, we may assume ρ is real and positive. From Theorem 2.1 we know that the differential dh has simple zeros at 0 and $\pm p$ and no poles at any other finite points. At the end e , it must have a triple pole (producing end-behavior like on Enneper's surface). The one-form dz has precisely these zeros and poles. Hence, $dh = cdz$ for some nonzero complex constant c , which by scaling we may assume to be unitary. We will now show that c is real. To see this, observe that along lifts of the real axis $\frac{dz}{\gamma}(\frac{\partial}{\partial t})$ is real as is $dz(\frac{\partial}{\partial t})$. But these curves are geodesic curvature lines, so by (2.13), c must be real. We may therefore choose $c = 1$ and conclude that

$$g = \rho\gamma \quad dh = dz, \quad (2.19)$$

where z is the projection defined above and $\rho > 0$ is to be determined.

All that remains is to choose $\rho > 0$ so that (2.8) is satisfied. Clearly, the third component has no periods because z is well-defined on the square torus. Because e lies on the intersection of two vertical symmetry planes, the real period of X must be orthogonal to both symmetry planes, and hence zero, for any closed curve around e .

To determine the value of ρ that makes X in (2.6) well-defined, we exploit the symmetry of the situation. Consider the curve on the square torus that is the lift of the interval $[-1, 0]$ in the z -plane, on which γ is positive. (It is always real). This curve must be in the vertical plane $x_2 = 0$, assuming we start the integration from $(0, 0)$. We require that

$$\operatorname{Re} \int_0^{-1} (g^{-1} - g) dh = 0,$$

or

$$0 = \operatorname{Re} \int_0^{-1} \left(\frac{1}{\rho\gamma} - \rho\gamma \right) dz = \rho^{-1} \int_{-1}^0 \sqrt{\frac{(t^2 - 1)}{t}} dt - \rho \int_{-1}^0 \sqrt{\frac{t}{(t^2 - 1)}} dt.$$

Both of the integrals are convergent and positive. Choosing ρ to be the positive square root of the ratio of these integrals insures that this way of evaluating $-p = X((-1, 0))$ places it on the x_3 -axis. It is evident that the choice of negative square roots will result in the same value of ρ , placing $X((1, 0))$ on the vertical axis. Thus X is well-defined on the torus and produces the genus-one Chen-Gackstatter example with the symmetry of Enneper's surface.

Remark 2.3 *i) The only complete minimal surfaces in \mathbb{R}^3 with total curvature -4π are the Catenoid and Enneper's surface (Osserman [58]). Recently Lopez [49] proved that the unique genus-one complete minimal surface with total curvature -8π is the Chen-Gackstatter example. This result was also contained in the thesis of D. Bloss [3].*

ii) The Chen-Gackstatter construction has been generalized in two different ways. First, one can add more handles. Chen and Gackstatter did this themselves in the genus-two case [11]. Computationally, higher genus examples were found by E. Thayer by a method that using a computer to solve the period problem (2.8). [69]. N. do Espírito-Santo generalized the method of proof in the genus two case to genus three [19]. Second, Karcher realized that the k th-order Enneper surface given by the Weierstrass data

$$g(z) = z^k, dh = g(z)dz, z \in C,$$

could also be used as a model for the end-behavior and symmetry of genus one examples. Thayer observed that the Chen-Gackstatter proof of existence extended to these higher order surfaces in the case of genus two. (See Remark 4.2.) He also computed higher genus examples (up to 45) with higher winding orders.

2.3 Embeddedness and finite total curvature: necessary conditions

We continue the general discussion of Section 2.1. Suppose $X: M \rightarrow \mathbb{R}^3$ has an embedded end at some $p \in \overline{M}_k - M$. Let $D - \{p\}$ be a punctured neighborhood of p on which X is one-to-one. We may assume, after a rotation if necessary, that the Gauss map, which extends to p , takes on the value $(0, 0, 1)$ at p . The two simplest examples are the plane and (either end of) the catenoid. Written as graphs over the (x_1, x_2) -plane, these ends have bounded and logarithmic growth respectively. This is in fact all that can happen.

Proposition 2.1 ([59] [66]) *Suppose $X: M \rightarrow \mathbb{R}^3$ has a complete embedded end $X: D - \{p\} \rightarrow \mathbb{R}^3$ of finite total curvature with the value of the extended Gauss map equal to $(0, 0, \pm 1)$ at p . Then outside of a compact set, $X(D - \{p\})$ is the graph (over the exterior of a bounded domain in the (x_1, x_2) -plane) with the following asymptotic behavior:*

$$x_3(x_1, x_2) = \alpha \log \rho + \beta + \rho^{-2}(\gamma_1 x_1 + \gamma_2 x_2) + \mathcal{O}(\rho^{-2}) \quad (2.20)$$

for $\rho = (x_1^2 + x_2^2)^{1/2}$. Moreover the first two components ϕ_1, ϕ_2 of Φ in (2.7) have poles of order two at p and have no residues, while the third component $dh = \phi_3$ is either regular (which happens if and only if $\alpha = 0$ in (2.20)) or has a simple pole.

Proof. We will prove this proposition in the case that the Gauss map at the embedded end is one-to-one. In this case we may assume, without loss of generality, that $p = 0$ and $g(z) = z$ on D . (In general we may assume $g(z) = z^k$ on D . The reader is invited to carry out a similar computation, showing that $\alpha = 0$ when $k > 1$.)

From (2.7) and (2.8) we have

$$\begin{aligned} x_1(z) &= \operatorname{Re} \int_{z_0}^z \phi_1 = \operatorname{Re} \int_{z_0}^z \frac{1}{2}(g^{-1} - g)dh = \operatorname{Re} \int_{z_0}^z (z^{-1} - z)dh \\ x_2(z) &= \operatorname{Re} \int_{z_0}^z \phi_2 = \operatorname{Re} \int_{z_0}^z \frac{1}{2}(g^{-1} + g)dh = \operatorname{Re} \int_{z_0}^z \frac{1}{2}(z^{-1} + z)dh \\ x_3(z) &= \operatorname{Re} \int_{z_0}^z dh \end{aligned}$$

It is straightforward to compute

$$x_1 - ix_2 = \int_{z_0}^z g^{-1}dh - \overline{\int_{z_0}^z gdh} = \int_{z_0}^z z^{-1}dh - \overline{\int_{z_0}^z zdh}.$$

We will now look more closely at dh . Writing

$$dh = \left(\sum_{j=1}^k c_{-j} z^{-j} + c_0 + zw_1(z) \right) dz,$$

$w_1(z)$ holomorphic, we observe that $x_1 - ix_2$ is not well-defined if $c_0 \neq 0$. The first integral in the expression for $x_1 - ix_2$ would produce a term $c_0 \ln z$, which is not a well-defined

function of z . For similar reasons (using the second integral) $c_{-2} = 0$. Knowing that $c_0 = 0$ allows us to conclude that dh has a pole at 0; otherwise the metric (2.9)

$$\frac{1}{2}(|g| + |g|^{-1})|dh| = \frac{1}{2}(|z| + |z|^{-1})|dh| = \frac{1}{2}(|z|^2 + 1)\frac{|dh|}{|z|}$$

is not complete at $|z| = 0$. We claim now that dh has a simple pole at $z = 0$. If not, let $c_{-k}z^{-k}, k \geq 3$, be the highest order term in the expansion. Then the dominant term in $\int zdh$ is $-(c_{-k}/k)z^{-k}, k \geq 3$. This means that on $|z| = \varepsilon$ small, $x_1 - ix_2(z)$ has self intersections. However, since we are assuming the end is embedded, this contradicts the fact that the normal is close to vertical near 0, so the projection of the image of a small circle $|z| = \varepsilon$ is one-to-one. Thus, $dh = (c_{-1}z^{-1} + zw_1(z))dz, w_1(z)$ holomorphic, as claimed. Because $x_3 = \operatorname{Re} \int_{z_0}^z dh$ is well-defined, c_{-1} must be real. Moreover, ϕ_1 and ϕ_2 have double poles without residues at $z = 0$.

We now can write

$$\begin{aligned} x_1 - ix_2 &= -c_{-1}/z + c_0 + zw_2(z) + \overline{zw_3(z)} \\ x_3 &= \operatorname{const} + c_1 \ln |z| + \mathcal{O}(|z|^2), \end{aligned}$$

$w_i(z)$ holomorphic. Because the end is complete, $c_1 \neq 0$. We compute

$$\rho^2 = |c_{-1}|^2/|z|^2 - 2\operatorname{Re}(\bar{c}_{-1}c_0/\bar{z}) + \operatorname{const} + \mathcal{O}(|z|)$$

and $\operatorname{Re}(x_1 - ix_2) \cdot \bar{c}_0$, which we express as follows:

$$\operatorname{Re}(c_{-1}/z)\bar{c}_0 = -\operatorname{Re}(x_1 - ix_2) \cdot \bar{c}_0 + |c_0|^2 + \mathcal{O}(|z|).$$

From the expression for ρ^2 one sees that $|z| = \mathcal{O}(\rho^{-1})$. Using this fact, and substitution of the second equation into the expression for ρ^2 gives, after rearrangement:

$$\begin{aligned} |c_{-1}|^2/|z|^2 &= \rho^2 + 2\operatorname{Re}(x_1 - ix_2 \cdot \bar{c}_0) + \operatorname{const} + \mathcal{O}(\rho^{-1}) \\ &= \rho^2(1 + \rho^{-2})(2\operatorname{Re}(x_1 - ix_2 \cdot c_0) + \mathcal{O}(\rho^2)). \end{aligned}$$

Taking the logarithm of both sides and using the fact that $\log(1 + a) = a + \mathcal{O}(a^2)$, we get

$$\ln |z| = \ln |c_1| - \ln \rho - \frac{\operatorname{Re}(x_1 - ix_2) \cdot c_0}{\rho^2} + \mathcal{O}(\rho^{-2})$$

substitution in the equation above for $x_3(z)$ yields:

$$x_3(z) = \operatorname{const} - c_1 \ln \rho - \rho^{-2}c_1 \operatorname{Re}(x_1 - ix_2) \cdot c_0 + \mathcal{O}(\rho^{-2}) \quad (2.20)'$$

which gives (2.20).

Definition 2.4 Consider an embedded end of a complete minimal surface of finite total curvature. The end is said to be a flat or planar end if $\alpha = 0$ in equation (2.20), and is a catenoid end otherwise. The constant α is called the logarithmic growth of the end.

Remark 2.4 From (2.20)':

- i) It is evident that an embedded finite total curvature end is asymptotic to an actual catenoid end ($h = \alpha \log r + \beta$) or to a plane ($h \equiv \beta$);
- ii) It also follows from equations (2.20)' that the logarithmic growth of a complete embedded end, parameterized on a punctured disk $D - \{p\}$, is equal to $-c_1$, where c_1 is the residue of dh at p . If the end is parameterized by the exterior of a disk ($p = \infty$) then the logarithmic growth is equal to the residue of dh at infinity;
- iii) In the proof of Proposition 2.1 we showed that if the extended Gauss map is unbranched at an embedded end and of finite total curvature, that end is a catenoid end. Since the only other geometric possibility is to have a flat end, it follows that branching of the extended Gauss map at an embedded end of finite total curvature is equivalent to the flatness of that end.

Proposition 2.2 (Gackstatter [24] Jorge-Meeks [42])

- i) All of the ends of M are embedded $\Leftrightarrow n(M) = r \Leftrightarrow$ equality holds in (2.14):

$$\int K dA = -4\pi(k + r - 1). \quad (2.21)$$

- ii) If no two ends of M intersect then, after a rotation, the Gauss map N satisfies $N(p_i) = (0, 0, \pm 1), i = 1, \dots, r$.

Remark 2.5 If $X: M \rightarrow \mathbb{R}^3$ is an embedding, then the conditions of Proposition 2.2 must hold. However, these conditions are not sufficient to imply that X is an embedding. See the discussion and examples presented in Section 5.

Proof i) follows immediately from Theorem 2.3 ii), formula (2.16) and the observation made after Definition 2.3 that $d_j = 1$ when an end is embedded. Statement ii) follows from Remark 2.4, because two ends that are of planar- or catenoid-type must intersect if their limit normals are not parallel.

Because formula (2.21) is important and we did not present the proof of Theorem 2.3 we will give a direct proof of it here. Formula (2.21) is clearly true for the flat plane ($k = 0, r = 1, K \equiv 0$). Hence we may assume $K < 0$ except at isolated points. M is by assumption a compact Riemann surface of genus k , from which r points, p_1, \dots, p_r , have been removed. Choose closed curves $\tilde{\beta}_j$ that bound small disks D_j centered at p_j . Since the ends are assumed to be embedded, Remark 2.4,i) implies that they are asymptotic to half-catenoids or planes. This means that a sequence of curves homotopic to $\tilde{\beta}_j$, whose lengths approach a minimum for all such curves, cannot diverge. Because $K < 0$ except at isolated points of M , a unique length-minimizing geodesic β_j exists in the homotopy class

of $\tilde{\beta}_j$ ([27]). Each geodesic bounds an end-representative that is topologically an annulus. By intersecting a planar end with a large sphere or a catenoid end with a plane orthogonal to the limit normal and sufficiently far out, we may approximate the annular end E_j by a finite annulus bounded on one side by the geodesic β_j and on the other by a closed curve c . Moreover the closed curve c can be chosen, according to Proposition 2.1 and formula (2.20), to be of the form $c_\rho(\theta)0 \leq \theta \leq 2\pi$, with

$$c_\rho(\theta) = (\rho \cos \theta, \rho \sin \theta, \alpha \log \rho + \rho^{-2}(\gamma_1 \cos \theta + \gamma_2 \sin \theta) + \mathcal{O}(\rho^{-2})).$$

Here $\rho > 0$ can be made as large as desired, and γ_1, γ_2 are fixed. Increasing ρ corresponds to pushing $c_\rho(\theta)$ farther out on the end; as $\rho \rightarrow \infty$, $c_\rho(\theta)$ diverges. It is straightforward to see from the equation for $c_\rho(\theta)$ that its total curvature converges quickly to 2π . Using the Gauss-Bonnet formula, we have

$$\int_{E_j} K dA \cong 2\pi\chi(E_j) - \int_{\beta_j} k_g ds - \int_c \kappa ds = - \int_c \kappa ds,$$

since the Euler characteristic of an annulus is zero and β_j is a geodesic. But since c is asymptotically a circle, we have shown that the total curvature of the end bounded by β_j is exactly -2π .

We next show that the β_j are pairwise disjoint. If two touch but do not cross, then of course they must coincide, which implies that S is the union of two ends with a common boundary. The surface S is an annulus ($g = 0, r = 2$) with total curvature -4π , so (2.21) is satisfied in this case. (By a result of Osserman [58], the only such surface is the catenoid.) If two β_j cross, then a segment of one together with a segment of the other bounds a region Σ of the surface. Since Σ is simultaneously inside of two ends it is easy to see that Σ is simply connected. This leads to a contradiction of the fact that each α_j is the unique geodesic in its homotopy class. (Alternatively, we could use the Gauss-Bonnet formula again to get a contradiction; the Euler characteristic of Σ is 1 and its boundary consists of two geodesics meeting at two vertices whose exterior angles are less than π in absolute value:

$$2\pi = 2\pi\chi(\Sigma) = \int_{\Sigma} K dA + \Sigma \text{ exterior angles} < \int_{\Sigma} K dA + 2\pi.$$

This clearly contradicts the fact that $K < 0$ almost everywhere on Σ .) Hence the β_j are disjoint.

Now remove the r ends from S by cutting along the geodesics α_j $j = 1, \dots, r$. This leaves a compact surface \widehat{M} of genus k , bounded by r closed geodesics. A third (and final) application of Gauss-Bonnet gives

$$2\pi(2 - 2k - r) = 2\pi\chi(\widehat{M}) = \int_{\widehat{M}} K dA.$$

Combining this with the previously established fact that each end has total curvature -2π gives

$$\int_M K dA = \int_{\widehat{M}} K dA + r(-2\pi) = -4\pi(k + r - 1),$$

which is (2.21). See [9] for more details.

If S is embedded and complete with finite total curvature, then we know by Proposition 2.1 that outside of a sufficiently large compact set of \mathbb{R}^3 , S is asymptotic to a finite number of half-catenoids and planes, which may be assumed to have the same vertical limit normal (Proposition 2.2 *ii*). If S is not a plane it must have *at least two* catenoid ends, one with positive, the other with negative logarithmic growth. This follows from the Halfspace Theorem (Theorem 2.4, which will be proved in Section 2.3.3). It states that a complete properly immersed minimal surface without boundary in \mathbb{R}^3 , which is not a flat plane, cannot lie in any halfspace. If S has only one catenoid end, it would be in a halfspace yet not be a plane, a contradiction.

Since we are assuming that S is connected, the fact that S is properly embedded implies that $\mathbb{R}^3 - S$ consists of precisely two components. Outside of a sufficiently large compact set, the ends of S are stacked and thus ordered from top to bottom. This also means that the limit normals $(0, 0 \pm 1)$ alternate from one end to the next and that the logarithmic growth rates are also ordered: If the ends correspond to points $\{p_1 \dots p_r\} \subset \overline{M}$, and α_j is the logarithmic growth rate of the j th end:

$$\alpha_1 \leq \alpha_2 \leq \dots \leq \alpha_r, \quad \alpha_1 \alpha_r < 0. \quad (2.22)$$

2.3.1 Flux

On a surface S , with boundary, let ν denote the outward-pointing unit-normal vector field on ∂S . For any vector field W that is C^1 with compact support on S , the divergence theorem states that

$$\int_S (\operatorname{div}_S W) dA = \int_{\partial S} \langle W, \nu \rangle ds.$$

If $S \subset \mathbb{R}^3$ and W is a vector field on \mathbb{R}^3 , we may restrict W to S and consider its tangential component $W^T := W - \langle W, N \rangle N$. Suppose W is a Killing field and S a minimal surface. Then we have $\operatorname{div}_S W^T = 0$, which can be seen as follows. For any tangent vector U , and any vector field: $W \in \mathbb{R}^3$

$$\begin{aligned} \nabla_U W^T &= [UW^T]^T = [UW - U(\langle W, N \rangle N)]^T \\ &= [UW]^T - \langle W, N \rangle DN(U). \end{aligned}$$

Since W is Killing, $\langle (UW)^t, U \rangle = \langle UW, U \rangle = 0$. Since S is minimal, $\operatorname{tr}_S DN = 0$. Hence $\operatorname{div}_S W^T = \operatorname{tr}_S \nabla W^T = \operatorname{tr}_S [DW]^T - \langle W, N \rangle \operatorname{tr}_S DN = 0$, as claimed.

The divergence theorem yields

$$\int_{\partial S} \langle W, \nu \rangle ds = 0,$$

since $\langle W^T, \nu \rangle = \langle W, \nu \rangle$. Thus the one-form $\omega(V) := \langle V, W \rangle$ is a closed form on S and for closed curves γ , $\int_\gamma \omega$ is an homology invariant.

Figure 2.3.1 Flux

Flux on a curve is computed by integrating the outward-pointing unit normal vector on a cycle. The total flux of the boundary of a minimal surface is zero.

A constant vector field T is a Killing field associated to translation in that direction. Since

$$\int_{\gamma} \langle T, \nu \rangle ds = \langle T, \int_{\gamma} \nu ds \rangle$$

and the left-hand side is a homology invariant for all constant vector fields T , so is $\int_{\gamma} \nu ds$.

Definition 2.5 *The flux along γ is the vector quantity*

$$Flux ([\gamma]) := \int_{\gamma} \nu(s) ds.$$

We remark that the flux can also be defined by the same equation for any closed curve $\gamma \in M$ using $\nu(s) = Rot_{\frac{\pi}{2}}(d\gamma/ds)$. This coincides with our definition of flux when γ is a boundary curve of M and is easily seen to be homology-invariant.

Proposition 2.3 *Suppose $X: M \rightarrow \mathbb{R}^3$ is a complete conformal minimal immersion given by $X = Re \int \Phi$ as in (2.6), (2.7). Suppose $\gamma = X(\tilde{\gamma}), \tilde{\gamma} \subset M$ is a closed curve. Let X^* denote the conjugate surface (See Definition 2.7), which is a (possibly multi-valued) conformal immersion $X^*: M \rightarrow \mathbb{R}^3$:*

$$i) \quad Flux ([\gamma]) = -Period_{\tilde{\gamma}} X^*;$$

ii) If $\tilde{\gamma}$ is in the homology class of a puncture, p ,

$$\text{Flux}([\gamma]) = -2\pi \text{Residue}_p \Phi;$$

iii) If $\tilde{\gamma}$ is in the homology class of a puncture p , which represents a vertical embedded end of finite total curvature,

$$\text{Flux}([\gamma]) = (0, 0, 2\pi\alpha), \quad (2.23)$$

where α is the logarithmic growth of the end. In particular the flux of a homology class representing a finite-total-curvature end is always vertical and is zero if and only if the end is flat;

iv) If $X(M)$ is an embedding with finite total curvature and logarithmic growth rates $\alpha_1, \dots, \alpha_r$ at the ends, then

$$\sum_1^r \alpha_j = 0. \quad (2.24)$$

Proof. A constant vector field T is the gradient, in \mathbb{R}^3 of a linear function f . The tangential component T^T of T restricted to $\mathcal{S} := X(M)$ is the (Riemannian) gradient of $f|_{\mathcal{S}}$, which is harmonic, according to (2.5), because X is a minimal immersion. Moreover, $\text{Rot}_{\frac{\pi}{2}} T^T = \text{grad } f^*$, where f^* is the locally defined harmonic conjugate of f . (See the discussion between (2.3) and (2.4).) Now we compute

$$\begin{aligned} T \cdot \text{Flux}([\gamma]) &= \int_{\gamma} \langle T^T, \nu \rangle ds = \int_{\gamma} \langle \text{Rot}_{\frac{\pi}{2}} T^T, \text{Rot}_{\frac{\pi}{2}} \nu \rangle ds \\ &= \int_{\gamma} \langle \text{grad } f^*, \frac{-d\gamma}{ds} \rangle ds = - \int_{\gamma} df^*. \end{aligned}$$

Since this is true for *any* constant vector field T in \mathbb{R}^3 , it follows that $\text{Flux}([\gamma]) = -\text{Period}_{\tilde{\gamma}} X^*$, which is i). From this, ii) follows immediately since

$$\int_{\tilde{\gamma}} \Phi = \text{Re} \int_{\tilde{\gamma}} \Phi + i \text{Im} \int_{\tilde{\gamma}} \Phi = \text{Period}_{\tilde{\gamma}} X + i \text{Period}_{\tilde{\gamma}} X^* = -i \text{Flux}([\gamma]). \quad (2.25)$$

To establish iii) we use Remark 2.4ii), together with statement ii) above and the fact that the third component of Φ is dh . We have that the third component of $\text{Flux}([\gamma])$, where $[\gamma]$ is the homotopy class of a puncture p , must equal $-2\pi \text{Residue}_p dh = 2\pi\alpha$, where α is the logarithmic growth of the end at p . According to Proposition 2.1, the first two components of Φ have no residues at p .

If $X(M)$ has finite total curvature then according to Theorem 2.2iii), dh is globally defined, with poles only at the ends $p_1 \dots p_r$. Hence, using statement iii)

$$0 = \sum_{i=1}^r \text{Residue}_{p_i} dh = 2\pi \left(\sum_{i=1}^r \alpha_i \right).$$

This establishes statement iv). □

Remark 2.6 *If we consider the normal component of the Killing field, $W^N =: w^N N$, where $w^N = \langle W, N \rangle$, a calculation shows that*

$$-\Delta_S w^N + 2K w^N = 0,$$

where K is the Gauss curvature of S . That is, $w^N N$ is a Jacobi-field. From a geometric point of view this is expected, since a Killing field is an infinitesimal isometry, and the Jacobi equation is derived from the formula for the second variation of area of a minimal surface. This is considered in more detail in Section 7.

2.3.2 Torque

Let $R_{\vec{u}}$ be the Killing field associated with counter-clockwise rotation about the axis $\ell_{\vec{u}}$ in the \vec{u} direction. From the identity $(U \wedge V) \cdot W = \det(U, V, W)$, for vectors U, V, W in \mathbb{R}^3 we have

$$(X \wedge \nu) \cdot \vec{u} = (\vec{u} \wedge X) \cdot \nu = R_{\vec{u}} \cdot \nu, \quad (2.26)$$

where X is the position vector of a minimal surface S and ν is the outward-pointing normal to a component γ of ∂S . Because $R_{\vec{u}}$ is a Killing field, $\int_{\gamma} R_{\vec{u}} \cdot \nu$ is a homology invariant and $\int_{\partial S} R_{\vec{u}} \cdot \nu = 0$. This motivates defining the *torque* of a closed curve γ on S as the vector-valued quantity.

Definition 2.6

$$Torque_0(\gamma) = \int_{\gamma} X \wedge \nu.$$

Torque was introduced by Kusner in [45, 46]. From (2.26) it follows that the component of torque in the \vec{u} direction is $\int_{\gamma} R_{\vec{u}} \cdot \nu$. If we move the origin from 0 to $W \in \mathbb{R}^3$ and let \hat{X} be the position vector measured from W , then $\hat{X} = X - W$ and we can compute

$$\begin{aligned} Torque_W(\gamma) &= Torque_0(\gamma) - W \wedge \int_{\gamma} \nu \\ &= Torque_0(\gamma) - W \wedge Flux(\gamma). \end{aligned} \quad (2.27)$$

It follows that if $Flux(\gamma) = 0$, the torque of γ does not depend upon the base-point of X . It also follows that the torque does not change if we move the base point parallel to the flux. If E is an embedded end we may speak of the torque of E ; it is the torque of a closed curve in the homology class of the ends. From Proposition 2.1 the following geometric proposition can be deduced.

Proposition 2.4 *If E is a vertical catenoid end the torque of E is a horizontal vector. If E is a vertical flat end at which the degree of the Gauss map is at least three, the torque of E vanishes.*

In the case of a vertical catenoid end, we know from Proposition 2.3iii) that the flux is a vertical vector of the form $(0, 0, 2\pi\alpha)$ where α is the logarithmic growth. We can conclude from (2.27) that there is a unique horizontal vector $W = (w_1, w_2, 0)$ for which the *Torque* of E is vertical. From Proposition 2.4, it follows that there is a unique vertical line on which the torque vanishes.

Definition 2.7 *For a vertical catenoid end E , the (unique) vertical line ℓ_E on which the torque vanishes is called the axis of E .*

The torque of E measured from a base point displaced by W from ℓ_E is

$$\begin{aligned} \text{Torque}_W(E) &= W \wedge (0, 0, 2\pi\alpha) \\ &= 2\pi\alpha(w_2, -w_1, 0). \end{aligned} \tag{2.28}$$

In the case of an embedded finite total curvature minimal surface with catenoid ends $E_1 \dots E_n$ and either no flat ends or all flat ends of order three or greater, the total torque of all the ends must sum to zero. For any base point let W_i , be the horizontal displacement of ℓ_{E_i} , $i = 1 \dots n$. Then

$$\begin{aligned} 0 &= \Sigma \text{Torque}_{-W_i}(E_i) = 2\pi \text{Rot}_{\frac{\pi}{2}}(\Sigma_{i=1}^n \alpha_i W_i), \quad \text{or} \\ \Sigma \alpha_i W_i &= 0. \end{aligned} \tag{2.29}$$

Remark 2.7 *If a surface has only two catenoid ends E_1, E_2 , and either no flat ends or all flat ends of order three or greater, then placing the base point on ℓ_{E_1} forces $W_1 = 0$ and by (2.29) we must have $W_2 = 0$. That is the catenoid ends E_1 and E_2 have the same axis. Pascal Romon [personal communication] points out that if there are three catenoid ends (and the same hypothesis about flat ends) their axes are coplanar. We may assume, without loss of generality that there are two ends with positive logarithmic growth. If we label these ends E_1 and E_2 and the third end E_3 and choose a base point on E_3 , then it follows from (2.29) that $W_1 = -\frac{\alpha_2}{\alpha_1} W_2$. Thus the axes ℓ_{E_1}, ℓ_{E_2} and ℓ_{E_3} are coplanar and, because $\frac{\alpha_2}{\alpha_1} < 0$, the axis ℓ_{E_3} with negative logarithmic growth lies between the two with positive growth. We note that on all known embedded examples, flat ends of order two do not occur. It is not known if they can occur.*

2.3.3 The Halfspace Theorem

We present here a proof of

Theorem 2.4 (The Halfspace Theorem for Minimal Surfaces, [40]) *A complete, properly immersed, nonplanar minimal surface in \mathbb{R}^3 is not contained in any halfspace.*

Our proof requires the use of the Maximum Principle for Minimal Surfaces: *If S_1 and S_2 are two connected minimal surfaces with a point p in common, near which S_1 lies on*

Figure 2.3.3 The vertically translated half-catenoid C
in the proof of Theorem 2.4.

one side of S_2 , then $S_1 = S_2$ or one is a subset of the other. We refer the reader to [17] for proofs. In the simple case that S_2 is a plane, the Maximum Principle states that a minimal surface $S = S_1$ that lies locally on one side of its tangent plane $T_p S = S_2$ must in fact be planar. This is easy to prove from the Weierstrass representation, Theorem 2.1. Without loss of generality, we may assume that $T_p S = \{x_3 = 0\}$ and that z is a local coordinate chart chosen so that $X(0) = p$. The Gauss map of S is vertical at p , so that $g(0) = \sigma \circ N(p) = 0, \infty$. Either g is constant, in which case S is flat and is a subset of $T_p S = \{x_3 = 0\}$, or 0 is an isolated pole or zero of g of order $k \geq 1$. In the latter case,

$$dh = z^k w(z) dz,$$

where $w(z)$ is holomorphic and $c = w(0) \neq 0$. Then

$$x_3(z) = \operatorname{Re} \int_0^z dh = \operatorname{Re} \int_0^z z^k w(z) dz = \operatorname{Re} \left(\frac{cz^{k+1}}{k+1} \right) + \mathcal{O}(|z|^{k+2}).$$

Clearly $x_3(z)$ changes sign $2(k+1)$ times in any neighborhood of $z = 0$. Hence S does not lie on one side of $T_p S = \{x_3 = 0\}$ near p .

Proof of Theorem 2.4. By assumption, S is contained in some halfspace H . The intersection, H^* , of the closed halfspaces containing S and having boundary parallel to ∂H is the smallest closed halfspace containing S . We will rotate and translate S so that $H^* = \{x_3 \geq 0\}$, and hence $\partial H^* = \{x_3 = 0\}$. If $S \cap \{x_3 = 0\} \neq \emptyset$, then the simple case of the Maximum Principle implies S is a plane. If S is not a plane $S \cap \{x_3 = 0\} = \emptyset$, and since S is properly immersed S has no limit points in $\{x_3 = 0\}$. This means each point of $\{x_3 = 0\}$ has positive distance from S .

Consider a catenoid with axis parallel to the x_3 -axis and waist circle in $\{x_3 = 0\}$. Let C denote the half-catenoid in $\{x_3 \leq 0\}$. Because x_3 is proper (and unbounded) on C it

follows that we can vertically translate C so that the waist circle is in $\{x_3 > 0\}$ but the *solid* half-catenoid inside C is disjoint from S . For convenience in the rest of the proof, we move our coordinate system so that $\partial H^* = \{x_3 = -\varepsilon\}$ and the waist-circle of C lies in $\{x_3 = 0\}$. Also, we will now think of C as the boundary for the solid half-catenoid i.e., we add to C the disk in $\{x_3 = 0\}$ bounded by the waist-circle.

Let tC denote the rescaling of C by a factor of $t > 0$. We will say that tC *lies below* S if S lies in the closure of the component of \mathbb{R}^3 above tC . (Similarly we may define what it means for a plane to lie below S .) For $t \leq 1$, the waist-circles of tC lie on C , so their distance from S is uniformly bounded away from S . If $S \cap tC \neq \emptyset$, then the Maximum Principle implies $S = tC$, and S being complete, must be a catenoid; this is a contradiction, since S lies in H^* . We may conclude that if tC is below S , then tC and S have no points in common. Furthermore, since $tC \cap H^*$ is compact and $S \subset H^*$, it follows that every tC lying below S actually has finite positive distance from S . In particular the set

$$T := \{t \in (0, 1] : tC \text{ is below } S\}$$

is open in $(0, 1]$. But if $\{t_n\} \subset T$ and $\lim_{n \rightarrow \infty} t_n = \tau > 0$, then τC is a half catenoid below S ; if not, some $t_n C$ must not be below S . Hence T is closed, too. This means that $T = (0, 1]$, so all tC are below S , $0 < t \leq 1$.

Note that the half-catenoids tC converge, as $t \rightarrow 0$, to the plane $\{x_3 = 0\}$ which is strictly inside $H^* = \{x_3 \geq -\varepsilon\}$. Since all the tC are below S , the limit plane $\{x_3 = 0\}$ is also below S . This contradicts the definition of H^* as the intersection of all the closed halfspaces containing S and having boundary parallel to ∂H , and this contradiction completes the proof. \square

Remark 2.8 *A slight modification of the same proof can be used to prove the following slight improvement of Theorem 2.4: Suppose S is a properly immersed nonplanar minimal surface with compact boundary ∂S . If H is a halfspace containing S , and H^* a smallest halfspace containing S , with ∂H^* parallel to ∂H , then $\partial H^* \cap \partial S \neq \emptyset$.*

2.4 Summary of the necessary conditions for existence of complete embedded minimal surfaces with finite total curvature

We conclude this section by gathering together the results of our discussion to formulate a sharpened version of Theorem 2.2 for *embedded* complete minimal surfaces of finite total curvature.

Proposition 2.5 *Let $X: M \rightarrow \mathbb{R}^3$ be a complete conformal minimal embedding with finite total curvature. Then:*

- i) M is conformally diffeomorphic to $\overline{M}_k - \{p_1, \dots, p_r\}$ where \overline{M}_k is a closed Riemann surface of genus k and p_1, \dots, p_r are points in \overline{M}_k , $r \geq 1$;*

- ii) X is proper. A punctured neighborhood of each p_i is mapped by X onto an end of $S = X(S)$ that is asymptotic to either a plane or a half-catenoid.
- iii) The Gauss map $N: M \rightarrow S^2$ extends to a meromorphic function on \overline{M}_k . All the normals on $\{p_1, \dots, p_r\}$ must be parallel and after a rotation if necessary we may assume that $N(p_i) = (0, 0 \pm 1)$, $i = 1, \dots, r$. The holomorphic one form dh , the complex differential of the height function, extends meromorphically to \overline{M}_k .
- iv) The total curvature of M is $-4\pi \cdot \text{degree } g = -4\pi(k+r-1)$, where $g = \sigma \circ N$, and $\sigma: S^2 \rightarrow \mathbb{C} \cup \{\infty\}$ is stereographic projection.
- v) The ends are naturally ordered by height from top to bottom. In this ordering, say $p_1 \dots p_r$, the unit vertical normals at the ends alternate and the logarithmic growth rates α_j are ordered from biggest to smallest. Furthermore,

$$\sum_{j=1}^r \alpha_j = 0. \quad (2.30)$$

If S is not a plane, $r \geq 2$ and $\alpha_1 \cdot \alpha_r < 0$.

- vi) At a catenoid end, g has a simple pole or zero, while at a flat end the pole or zero of g , has higher order. The height differential dh has a simple pole at a catenoid end. At a planar end where g has a zero or pole of order m , dh has a zero of order $|m| - 2 \geq 0$. The logarithmic growth rate is equal to minus the residue of dh at the puncture corresponding to the end.

To construct a complete embedded minimal surface of finite total curvature with the Enneper-Weierstrass-Riemann representation ((2.6) (2.7) of Theorem 2.1), Proposition 2.5 dictates necessary conditions on the choice of \overline{M}_k , $\{p_1 \dots p_r\} \subset \overline{M}_k$, $g = \sigma \circ N$ and dh . However, in order for (2.7) to be single-valued on $M_k = \overline{M}_k - \{p_1 \dots p_r\}$, it is necessary that (2.8) be satisfied:

$$\text{Period}_\alpha(\Phi) = \text{Re} \int_\alpha \left((g^{-1} - g) \frac{dh}{2}, i(g^{-1} + g) \frac{dh}{2}, dh \right) = 0 \quad (2.31)$$

for all closed curves $\alpha \subset M_k$. (See Figure 3.0.0.) If Statement iii) of Proposition 2.5 is satisfied, the resulting surface must have embedded parallel ends, which implies that outside of some compact set in \mathbb{R}^3 , $X(M_k)$ is embedded. However, X may not be an embedding. See the examples in Section 5.

3 Examples with restricted topology: existence and rigidity

In this section, we present the few uniqueness results that are presently known. They take the following form: if the topological type is restricted or some other geometric property is specified, then the surface is uniquely determined. (See also Theorem 3.4, below.)

Theorem 3.1 *Let S be a complete embedded minimal surface of finite total curvature.*

- 1) *If S has one end, it is a flat plane;*
- 2) *(The Lopez-Ros punctured sphere Theorem) [51]. If S has genus zero, it is the catenoid or the flat plane;*
- 3) *(The R. Schoen catenoid characterization) [66]. If S has two ends, it is the catenoid;*
- 4) *If S has three ends and a symmetry group of order at least $4(k+4)$, where k is the genus, then S is the surface described in Theorem 3.2.*
- 5) *(Costa's thrice-punctured torus Theorem) [16]. If S has genus one and three ends, it is one of the surfaces $M_{1,x}$ in Theorem 3.3.*

Statement 1) follows from Theorem 2.4 or Proposition 2.5 v). Statement 4) is proved in [36]. We will give a full proof of 2) in Section 3.1.

Corollary 3.1 *The only complete embedded minimal surfaces with $\int K dA \geq -8\pi$ are the plane and the catenoid (with total curvature 0 and -4π respectively).*

Proof. From Proposition 2.2, it follows that the total curvature is an integer multiple of -4π . If the total curvature is zero, we must have the plane. If it is -4π , then by a result of Osserman [58] it must be the catenoid. (See also Lemma 3.1 in Section 3.1.)

We need to show that total curvature -8π is not possible. In that case we would have, by Proposition 2.2, i), that $k+r=3$. But the various cases of Theorem 3.1 exclude this possibility. \square

Proposition 3.1 *Let $X: M \rightarrow \mathbb{R}^3$ be a connected complete minimal immersion with finite total curvature. Let $n(M)$ be as in Definition 2.3. Then for any $p \in \mathbb{R}^3$, the number of points in $X^{-1}(p)$ is at most $n(M) - 1$, with the sole exception of the case when $X(M)$ is the flat plane.*

Corollary 3.2 *Let $X: M \rightarrow \mathbb{R}^3$ be as in Theorem 3.1. If $n(M) = 2$, $X(M)$ is the catenoid.*

Proof of Proposition 3.1. We will use a version of the monotonicity formula to prove this proposition: *If p is a point on a complete immersed minimal surface and $A(r)$ is the area of the minimal surface inside a Euclidean ball of radius r , then $A(r)/\pi r^2$ is a nondecreasing function of r , which is strictly increasing unless the minimal surface is a collection of planes.*

Suppose that $p \in X(M)$ and $X^{-1}(p)$ contains n points. Assume that $X(M)$ is not a plane. (It cannot be a collection of planes because we assume that $X(M)$ is connected.)

Figure 3.0.0 Attempts to produce nonexistent minimal surfaces.

Left: According to the Lopez-Ros punctured sphere theorem (Theorem 3.1,2)), a complete embedded minimal surface with genus zero and finite total curvature is either the plane or the catenoid. Nonetheless, it is possible to write down Weierstrass data that would produce such an example, provided the period problem (2.8) were solvable (which it is not). One can do this with a high degree of symmetry. Under the assumptions that the desired example has three parallel vertical ends –top and bottom catenoidal, and the middle one flat– one vertical plane of symmetry and a horizontal line diverging into the flat end, the resulting Weierstrass data is

$$g = r(z - 1)(z + 1)^{-1}, \quad dh = dz$$

on $C' - \{\pm 1\}$ with $r > 0$. The gap in the image indicates the failure of the Weierstrass-Enneper mapping to be single-valued. The value of r chosen here is 1.

Right: According to the Schoen catenoid characterization (Theorem 3.1,3)) the catenoid is the only complete embedded minimal surface with two ends and finite total curvature. Nevertheless, we can find symmetric Weierstrass data that fails only in that the period problem is not solvable. Under the assumptions that the desired example has genus one, two vertical planes of reflective symmetry and one horizontal plane of reflective symmetry, the Weierstrass data is $\{g, dh\}$, where

$$g^2 = (1 - z)(r - z)((1 + z)(r + z))^{-1}, \quad dh = (z^2 - 1)^{-1}dz,$$

$r > 1$, on the rectangular torus determined by the equation relating g and z . The image looks like a catenoid through which someone has tried to drill a tunnel from both sides. The tunnels don't meet up as one sees here (for $r = 2.5$). By (Theorem 3.1,3)), it is futile to try other values of r .

Hence, for small r , $A(r)/\pi r^2 > n$. But as a consequence of Theorem 2.3, $A(r)/\pi r^2$ converges to $n(M)$ as $r \rightarrow \infty$. From monotonicity we conclude that $n < n(M)$. If $X(M)$ is a flat plane $n = 1 = n(M)$. \square

Proof of Corollary 3.2. It follows from Proposition 3.1 that X is an embedding and since $n(M) = 2$, it is not the plane. It must therefore have two embedded ends. From Theorem 3.1.3, it must be the catenoid. \square

According to Corollary 3.1 next smallest possible total curvature for an embedded example is -12π ; that is, $k + r = 4$. By Theorem 3.1, this can only happen when $k = 1$ and $r = 3$, and in fact it does.

Theorem 3.2 ([35, 36]) *For every $k \geq 2$, there exists a complete properly embedded minimal surface of genus $k - 1$ with three annular ends. After suitable rotation and translation, the example of genus $k - 1$, which we will call M_k , has the following properties:*

1. M_k has one flat end between its top and bottom catenoid ends. The flat end is asymptotic to the (x_1, x_2) -plane.
2. M_k intersects the (x_1, x_2) -plane in k straight lines, which meet at equal angles at the origin. Removal of the k lines disconnects M_k . What remains is, topologically, the union of two open annuli;
3. The intersection of M_k with any plane parallel (but not equal) to the (x_1, x_2) -plane is a single Jordan curve;
4. The symmetry group of M_k is the dihedral group with $4k$ elements generated by reflection in k vertical planes of symmetry meeting in the x_3 -axis, and rotation about one of the lines on the surface in the (x_1, x_2) -plane. Removal of the intersection of M_k with the (x_1, x_2) -plane and the vertical symmetry planes disconnects M_k into $4k$ congruent pieces, each a graph.
5. M_k is the unique properly embedded minimal surface of genus $k - 1$ with three ends, finite total curvature, and a symmetry group containing $4k$ or more elements.

As will follow from our presentation in Section 4: The Riemann surface \overline{M}_k is given by

$$w^k = -\left(\frac{1}{2}\right)z^{k-1}(z-1)(z+1). \quad (3.1)$$

The catenoid ends are located at $(z, w) = (\pm 1, 0)$, and the flat end is at (∞, ∞) . The Gauss map and the complex differential of the height function are given by

$$g = \rho w, \quad dh = \frac{dz}{(z-1)(z+1)} \quad (3.2)$$

Figure 3.0.1 Costa's surface (upper left) and a few members of the family of her deformations, as described in Theorem 3.3.

where ρ is a constant determined by the necessity of satisfying the period condition (2.6) in Proposition 2.5. The symmetry of the surface is used to show that there is only one period condition. (See Figure 3.1 for pictures of the M_k .)

Remark 3.1 *The example with $k = 2$ was found by Celso Costa in 1982. He proved it was complete and had embedded ends of the specified type. See Costa [14, 16] and Hoffman [28, 29].*

The surfaces M_k each lie in a family of embedded minimal surfaces.

Theorem 3.3 ([30]. See Section 4) *For every $k \geq 2$, there exists a one parameter family, $M_{k,x}, x \geq 1$, of embedded minimal surfaces of genus $k - 1$ and finite total curvature. The surfaces $M_{k,1}$ are precisely the surfaces M_k of Theorem 3.2. The surfaces $M_{k,x}, x > 1$ have all three ends of catenoid type and a symmetry group generated by k vertical planes of reflectional symmetry. The Riemann surface $\overline{M}_{k,x}$ is given by*

$$w^k = -c_x z^{k-1}(z - x)(z + x^{-1}), \quad c_x = (x + x^{-1})^{-1}. \quad (3.3)$$

Figure 3.0.2 Higher-genus embedded minimal surfaces.

The surfaces in the column on the left are the surfaces $M_k, k = 3, 4, 6$ described in Theorem 3.2, with genus 2 (top), 3 (middle) and 5 (bottom). The middle end in each of these surfaces is flat. In each row, the other two surfaces are deformations of the left-most one. These are the surfaces described in Theorem 3.3. Their middle ends are catenoidal.

The catenoid ends are located at $(z, w) = (x, 0), (-x^{-1}, 0)$ and (∞, ∞) . The Gauss map and differential of the height function are given by

$$g = \frac{\rho w}{mz + 1}, \quad dh = \frac{z(m + z^{-1})dz}{(z - x)(z + x^{-1})}, \quad (3.4)$$

where ρ and m constants determined by the period conditions. (When $x = 1, m(1) = 0$, and (3.3) and (3.4) give (3.1) and (3.2) for the surfaces $M_k = M_{k,1}$.)

Remark 3.2 *The surfaces $M_{k,1}$ of Theorem 3.3 are the surfaces M_k of Theorem 3.2. In Section 4, we find it more convenient to work with the function $u = z/w$ instead of w . The formulae (3.3) and (3.4) are presented (e.g. in (4.12)) in terms of u and z . See (4.4) where we perform the conversion, into a (w, z) expression, of the Riemann surface equation (3.3).*

In Section 5, we will discuss examples with more than three ends, some other results about the structure of the “space” of embedded finite total curvature examples, and present some questions and conjectures. In Section 4 we describe in detail the ideas behind the construction of the surfaces $M_{k,x}$ of Theorem 3.3. As we pointed out in the introduction, these are the only higher-genus examples that have been fully analyzed.

3.1 Complete embedded minimal surfaces of finite total curvature and genus zero: the Lopez-Ros theorem

In this section we prove the second statement of Theorem 3.1, which is Theorem 3.5 below. Our presentation follows that of Perez and Ros [60], whose approach yields the following theorem.

Theorem 3.4 *Let S be a complete embedded minimal surface with finite total curvature and vertical flux. Then S is the catenoid or the plane.*

Recall that for a closed curve $\gamma \subset S$ its flux is defined in Section 2.3.1 as

$$\text{Flux}([\gamma]) = \int_{\gamma} \text{Rot}_{\frac{\pi}{2}}\left(\frac{d\gamma}{ds}\right) ds.$$

$\text{Flux}([\gamma])$ depends only on the homology class of γ . We say that S has *vertical flux* provided $\text{Flux}([\gamma])$ is a vertical vector for all closed $\gamma \subset S$.

As usual, we may realize S by a conformal minimal embedding $X: M \rightarrow \mathbb{R}^3$, where $M = \overline{M} - \{p_1 \dots p_r\}$, and rotate S in \mathbb{R}^3 so that $g = 0$, or ∞ at the ends $\{p_1 \dots p_r\}$.

If S is complete and embedded, with genus zero, the only closed curves one has to consider are those associated to the ends $\{p_1 \dots p_r\}$. In Proposition 2.3ii) we established that the flux of the homology class of p_j is equal to the vertical vector $(0, 0, 2\pi\alpha_j)$, where α_j is the logarithmic growth of the end at p_j . Hence we have as an immediate consequence of Theorem 3.4:

Theorem 3.5 ([51]) *Let S be a complete minimal surface with genus zero and finite total curvature. Then S is the catenoid or the plane.*

The proof of Theorem 3.4 exploits what has come to be called the Lopez-Ros deformation. If $\{g, dh\}$ is the Weierstrass data of S , we define on \overline{M} the data

$$g_{\lambda} = \lambda g \quad dh_{\lambda} = dh, \quad (3.5)$$

for any $\lambda > 0$. Notice that the zeros and poles of $\{g_{\lambda}, dh_{\lambda}\}$ are the same as these of $\{g, dh\}$. The deformation we consider is given by the Weierstrass representation (2.6) (2.7).

$$X_{\lambda} := \text{Re} \int \Phi_{\lambda} := \text{Re} \int \left(\frac{1}{2}(\lambda^{-1}g^{-1} - \lambda g), \frac{i}{2}(\lambda^{-1}g^{-1} + \lambda g), 1 \right) dh. \quad (3.6)$$

Note that X_λ may be multivalued. The metric and curvature of X_λ are given by (2.9) (2.10) as

$$\begin{aligned} ds_\lambda &= (\lambda^{-1}|g|^{-1} + \lambda|g|)|dh| \\ K_\lambda &= \frac{-16}{(\lambda^{-1}|g|^{-1} + \lambda|g|)^4} \frac{|dg|}{|dh|}. \end{aligned} \tag{3.7}$$

We will prove three propositions about the Lopez-Ros deformation that allow us to prove Theorem 3.4. The propositions are stated here and then the proof of Theorem 3.4 is presented. The proofs of the propositions follow.

Proposition 3.2 *Let $X: M \rightarrow \mathbb{R}^3$ be a conformal minimal immersion:*

- i) X is complete if and only if X_λ is complete for all $\lambda > 0$;*
- ii) If X_λ is single-valued, the total curvature of X is finite if and only if the total curvature of X_λ is finite for all $\lambda > 0$;*
- iii) The immersions X_λ are single-valued for all $\lambda > 0$ if and only if X has vertical flux.*

Proposition 3.3 *Let $X: M \rightarrow \mathbb{R}^3$ be a complete embedded minimal surface of finite total curvature, for which X_λ is single-valued for all $\lambda > 0$. Then X_λ is an embedding for all $\lambda > 0$.*

Proposition 3.4 *Suppose $X: M \rightarrow \mathbb{R}^3$ is a conformal minimal immersion.*

- i) If for some $p \in M$, $N(p) = (0, 0 \pm 1)$ (i.e., $g(p) = 0$ or $g(p) = \infty$) then, on every neighborhood of p , X_λ is not an embedding, for λ sufficiently large.*
- ii) Suppose $X: M = \{z | 0 < |z| < \varepsilon\} \rightarrow \mathbb{R}^3$ is a conformal embedding representing a flat end. Then X_λ is not an embedding, for λ sufficiently large, unless X represents the end of a flat plane.*

Proof of Theorem 3.4. Let $X: M \rightarrow \mathbb{R}^3$ be a conformal minimal embedding with $S = X(M)$. The vertical flux condition guarantees that the Lopez-Ros deformation X_λ defined by (3.5) (3.6) is single-valued for all $\lambda > 0$ (Proposition 3.2 above). Because X is a complete embedding with finite total curvature, the same is true for X_λ , $\lambda > 0$ (Propositions 3.2, and 3.3 above). But this means that S has no vertical points (points where $g = 0, \infty$) and no flat ends, unless S is itself a flat plane (Proposition 3.4 above).

Assume that S is not a plane.

Because S is embedded and has finite total curvature, $M = \overline{M} - \{p_1 \dots p_r\}$, where \overline{M} is compact. The ends at $p \in \{p_1, \dots, p_r\}$ are either planar or catenoid-type. The zeros of dh can occur only at vertical points of g or at flat ends, while dh has simple poles at the

catenoid ends. (See Proposition 2.5 in Section 2.3.4.) From the previous paragraph we can conclude that dh has no zeros. (Because S is not a plane, $dh \neq 0$.) Thus $X(\overline{M}) = r > 0$, which implies that \overline{M} is the sphere and $r = 2$. We may conclude from Lemma 3.1 that S is the catenoid. \square

Lemma 3.1 *Suppose S is a complete embedded minimal surface with total curvature -4π and two ends. Then S is a catenoid.*

Proof of Lemma 3.1. Because $g = 0, \infty$ at the ends, and the values must alternate (see Proposition 2.5), we may assume, without loss of generality, that $\overline{M} - \{p_1, p_2\} = \mathbb{C} - \{0\}$ and $g(0) = 0, g(\infty) = \infty$. Since g has no other poles or zeros, it follows that $g(z) = az, a \neq 0$. Without loss of generality, we may assume that a is real, because rotation of S about a vertical axis changes g by a unitary multiplicative factor. Because dh has a simple pole at 0 and at ∞ and no other zeros or poles, $dh = cdz/z$ for some $c \neq 0$. Letting $\zeta = az$ we have $g(\zeta) = \zeta$ and $dh = cd\zeta/\zeta$. Clearly, dh has a real period on $|\zeta| = 1$ if c is not real. Thus our surface is the catenoid as described in Section 2. (See Figure 2.0.) \square

We now present proofs of Propositions.

Proof of Proposition 3.2. From (3.7) it is evident that for each $\lambda > 0$

$$\underline{c} ds \leq ds_\lambda \leq \overline{c} ds$$

$$\overline{c}^{-4} |K| \leq |K_\lambda| \leq \underline{c}^{-4} |K|, \text{ where}$$

$\overline{c} = \max\{\lambda^{-1}, \lambda\}$ and $\underline{c} = \min\{\lambda^{-1}, \lambda\}$. From these inequalities, statements i) and ii) follow directly. To prove statement iii), we use (2.25) from Section 2.3.1. For any closed curve $\gamma = X(\tilde{\gamma}), \tilde{\gamma} \subset M$, we have

$$\int_{\tilde{\gamma}} \Phi = \operatorname{Re} \int_{\tilde{\gamma}} \Phi + i \operatorname{Im} \int_{\tilde{\gamma}} \Phi = \operatorname{Period}_\gamma(X) - i \operatorname{Flux}_X([\gamma]) = -i \operatorname{Flux}_X([\gamma]).$$

From this identity it is evident that the vertical flux condition is equivalent to the exactness of ϕ_1 and ϕ_2 . Since $\phi_1 - i\phi_2 = g^{-1}dh$ and $-(\phi_1 + i\phi_2) = gdh$, exactness of ϕ_1 and ϕ_2 is equivalent to the exactness of $g^{-1}dh$ and gdh . But from (3.6), it is clear that $\operatorname{Period}_\gamma(X_\lambda) = \operatorname{Re} \int_{\tilde{\gamma}} \Phi_\lambda = 0$, for all $\lambda > 0$ and all $[\gamma]$, if and only if $g^{-1}dh$ and gdh are exact. \square

Proof of Proposition 3.3. Since the Lopez-Ros deformation (3.5) (3.6) changes neither zeros nor poles of g and dh , the end types of X_λ at the ends p_j of $M = \overline{M} - \{p_1 \dots p_r\}$ are the same for all $\lambda > 0$. That is, flat ends remain flat, and catenoid-type ends remain catenoidally catenoidal; even more, their logarithmic growth rates $\alpha = -\operatorname{Residue}_p dh$ are independent of λ .

Fix $\lambda_0 > 0$ and consider two distinct ends $p_i, p_j, i \neq j$. Choose disjoint neighborhoods D_i, D_j of p_i and p_j so that $X(D_i)$ and $X(D_j)$ are end-representatives. Suppose X_{λ_0} is an

embedding. If these ends have different logarithmic growths, i.e., if $\alpha_i \neq \alpha_j$, the distance function from $X(D_i)$ to $X(D_j)$ is not only nonzero but unbounded. This is because the ends are asymptotic to catenoid ends (or to a plane if one of the α 's is zero) with different logarithmic growth rates. Since these growth rates are independent of λ , it follows that $X_\lambda(D_i) \cap X_\lambda(D_j) = \emptyset$ for λ sufficiently close to λ_0 . In case $\alpha_i = \alpha_j$ and $X_{\lambda_0}(D_i)$ and $X_{\lambda_0}(D_j)$ are asymptotic to ends with the same growth rate, we appeal to the Maximum Principle at Infinity proved in [52], which states that the distance between these embedded annular ends is finite; i.e., they are *not* asymptotic at infinity. Thus, again, $X_\lambda(D_i) \cap X_\lambda(D_j) = \emptyset$ for λ sufficiently close to λ_0 . We may conclude that, for λ sufficiently close to λ_0 , X_λ is an embedding. Stated differently, what we have proved is that

$$L := \{\lambda > 0 \mid X_\lambda \text{ is injective}\}$$

is an open set. It is nonempty since $1 \in L$.

We now wish to show that L is closed. Suppose $\{\lambda_k\}$ is a sequence in L that converges to λ_0 , and suppose X_{λ_0} is *not* an embedding; i.e., $X_{\lambda_0}(q_1) = X_{\lambda_0}(q_2)$ for some points $q_1, q_2 \in M, q_1 \neq q_2$. Because X_{λ_k} converges, uniformly on compact subsets of M , to X_{λ_0} it follows that some neighborhood \mathcal{O}_1 of q_1 is mapped by X_{λ_0} to be on one side of the image of some neighborhood \mathcal{O}_2 of q_2 . By the Maximum Principle, stated in Section 2.3.3, $X_{\lambda_0}(\mathcal{O}_1) = X_{\lambda_0}(\mathcal{O}_2)$ and analyticity, $X_{\lambda_0}: M \rightarrow \mathbb{R}^3$ is a finite covering map, whose image $\mathcal{S} = X_{\lambda_0}(M)$ is an embedded minimal surface of finite total curvature in \mathbb{R}^3 .

Using the arguments presented above, we may assert that any two distinct ends of \mathcal{S} stay a bounded distance one from the other. This means that \mathcal{S} has an embedded tubular neighborhood \mathcal{U} of some radius $\varepsilon > 0$. For sufficiently large k , $X_{\lambda_k}(M) \subset \mathcal{U}$. Because the behavior of X_{λ_k} is completely controlled at the ends and $g_{\lambda_k} = \lambda_k g_k \rightarrow \lambda_0 g_k = g_{\lambda_0}$, the projection $\pi: \mathcal{U} \rightarrow \mathcal{S}$, when composed with X_{λ_k} , results in a finite covering map $\pi \circ X_{\lambda_k}$, for k sufficiently large. Because X_{λ_k} is one-to-one, the distance function in each fiber of \mathcal{S} separates points in the sheets of each covering map $\pi \circ X_{\lambda_k}$. The distance to \mathcal{S} is a continuous function on \mathcal{U} and hence the covering is single-sheeted. Since $\pi \circ X_{\lambda_k}$ converges uniformly, on compact subsets of M , to $\pi \circ X_{\lambda_0} = X_{\lambda_0}$ it follows that X_{λ_0} is in fact single-sheeted, i.e., an embedding, a contradiction. Hence $\lambda_0 \in L$. Thus we have shown that L is closed. Because L is also open and nonempty, $L = (0, \infty)$. Thus X_λ is an embedding for all $\lambda > 0$. \square

Proof of Proposition 3.4. Suppose q is a point where the Gauss map is vertical. Without loss of generality, we may assume that $g(q) = 0$. Choose conformal coordinates near q so that q corresponds to $z = 0$ and $g(z) = z^k$ for some $k > 0$ on $D_r(0) := \{|z| < r\}$. Because we are assuming regularity, dh must have a zero of order k at 0, i.e.,

$$g(z) = z^k \quad dh = z^k(a + zf(z))dz,$$

Figure 3.1 The limit surfaces in the proof of Proposition 3.4

Left: The k -Enneper surface given by (3.9), with $k = 2$.
 Right: The limit surface given by (3.10), with $k = 2$. There is one horizontal flat end and one immersed end with winding number equal to three.

where a is some nonzero complex constant and $f(z)$ is holomorphic on $D_r(0)$. For each $\lambda > 0$, the change of variables $z = \lambda^{-\frac{1}{k}}\zeta$ from $D_{\lambda^{\frac{1}{k}}r}(0)$ to $D_r(0)$ allows us to express g_λ and dh_λ as follows:

$$g_\lambda(z(\zeta)) = \zeta^k \quad dh_\lambda = dh = \lambda^{-(1+\frac{1}{k})}\zeta^k(a + \lambda^{-\frac{1}{k}}\zeta\dot{f}(\lambda^{-\frac{1}{k}}\zeta))dz \quad (3.8)$$

on $D_{\lambda^{\frac{1}{k}}r}(0)$.

Rescale dh_λ by a factor of $\lambda^{(1+\frac{1}{k})}$ so that $dh_\lambda = \zeta^k(a + \lambda^{-\frac{1}{k}}\zeta\dot{f}(\lambda^{-\frac{1}{k}}\zeta))dz$. The rescaled data produce immersions, which we will also call X_λ , that converge uniformly, on compact subsets of $\mathbb{C} = \lim_{\lambda \rightarrow \infty} D_{\lambda^{\frac{1}{k}}r}(0)$, to the immersion produced by the Weierstrass data

$$\widehat{g} = \zeta^k \quad \widehat{dh} = a\zeta^k dz. \quad (3.9)$$

We will call this immersion $\widehat{X}: \mathbb{C} \rightarrow \mathbb{R}^3$. The limit surface is a rescaled version of the k -Enneper surface defined and discussed in Remark 2.3ii). This surface is clearly not *embedded*. (See Figure 3.1.1.) Since rescaling does not create or destroy self-intersections, it follows that X_λ is not injective for λ sufficiently large. This completes the proof of the first statement of the proposition.

We now prove the second statement. Because a planar end has zero flux, X_λ is well-defined for all $\lambda > 0$, according to Proposition 3.2. Because it has finite total curvature, the Weierstrass data extends to zero meromorphically. Without loss of generality, we may assume that the limit normal at the end is $(0, 0, -1)$, so $g(0) = 0$. Because the end is flat,

the zero of g at $z = 0$ has order $k \geq 2$ and dh has a zero of order $k - 2$ at $z = 0$. After a change of variables, if necessary, we may assume that

$$g(z) = z^k \quad dh = z^k (az^{-2} + f(z))dz,$$

with $a \neq 0$ and $f(z)$ holomorphic on some disk $D_r(0) = \{|z| < r\}$. As in the proof of the first part of the proposition, we change variables by $z = \lambda^{-\frac{1}{k}}\zeta$ and rescale – this time by $\lambda^{1-\frac{1}{k}}$ – so that

$$g_\lambda(\zeta) = \zeta^k \quad dh = \zeta^k (a\zeta^{-2} + \lambda^{-\frac{2}{k}}f(\lambda^{-\frac{1}{k}}\zeta))d\zeta$$

on $D'_{\lambda^{\frac{1}{k}}r}(0)$. The associated immersions converge, uniformly on compact subsets of $\mathbb{R}^3 - \{0\} = \lim_{\lambda \rightarrow \infty} D'_{\lambda^{\frac{1}{k}}r}(0)$, to an immersion $\widehat{X}: \mathbb{R}^3 - \{0\} \rightarrow \mathbb{R}^3$ given the Weierstrass data

$$\widehat{g}(\zeta) = \zeta^k \quad \widehat{dh} = a\zeta^{k-2}d\zeta. \quad (3.10)$$

This immersion is complete and has a flat end at 0. However it has a nonflat end at infinity, and that end is not embedded. It follows that X_λ is not one-to-one for $\lambda > 0$ sufficiently large. \square

4 Construction of the deformation family with three ends

We wish to construct complete embedded minimal surfaces of genus $k - 1, k \geq 2$ with three catenoid ends and k vertical planes of symmetry that intersect in a common vertical line. They will be the surfaces $M_{k,x}$ of Theorem 3.3. These surfaces should be deformations of the examples M_k of Costa ($k = 2$) and Hoffman-Meeks [35, 36] ($k \geq 3$), given in Theorem 3.2, which have two catenoid ends and a middle flat end. There will be, for each genus, a one parameter family of such deformations. We begin by locating the zeros and poles of $g = \sigma \circ N$, the stereographic projection of the Gauss map. We will then show that the conformal type of the example is quite restricted.

Assume we have such a surface S for some fixed $k \geq 2$. Recall from Theorem 2.2 that the conformal type of S is that of a compact Riemann surface punctured in three points (one for each end). Each of these ends is fixed by each of the reflections. Thus the closure of the fixed point sets of these reflective isometries contain each of the puncture points. We express this by saying that: “Each of the k symmetry planes passes through each of the three punctures.”

Because the surface is properly embedded, it divides \mathbb{R}^3 into two regions. The vertical line of intersection of the symmetry planes may be taken to be the x_3 -axis. The symmetry planes make equal angles of π/k , otherwise the surface would have more symmetry planes by Schwarz reflection. The top and bottom ends must be catenoid-type. (This follows from Proposition 2.5. See also the discussion after Theorem 2.4.) Since these ends have the same limit-normals (because there are an odd number of ends and their normals must alternate

between $(0, 0, +1)$ and $(0, 0, -1)$) it follows that, for $x_3 > 0$ sufficiently large, the x_3 -axis is in one component of $\mathbb{R}^3 - S$, and in the other component for $x_3 < 0$, sufficiently negative. Thus the x_3 -axis passes through the surface an odd number of times.

We will show that it passes through *exactly once*. The reflections in vertical planes generate a cyclic rotation group of symmetries, of order k , about the x_3 -axis. Because S is embedded, at any point p of intersection of S and the x_3 -axis, S must have a vertical normal vector. The order of the zero or pole of g at p is one less than the number of curves in $S \cap T_p M$. If S has a rotational symmetry of order k this means that the zero or pole has order at least $k - 1$. On the other hand, it follows from Proposition 2.2 or Proposition 2.4 that

$$\text{degree } g = \text{genus } S + 2 = k + 1. \quad (4.1)$$

If there is more than one axis point, then the topmost and the bottommost points have the same orientation as the extreme catenoid ends. Orient S so that $g \circ N$ has zeros at these points. Then the number of zeros of g is at least $2(k - 1) + 2$, which is strictly greater than $k + 1$ for $k \geq 2$. This contradicts (4.1). Hence there is a unique axial point where g has a zero of order at least $k - 1$. But since g has a zero at one of the extreme catenoid ends this makes $k + 1$ zeros. We can conclude from (4.1) that the axial point is a zero of g of order exactly $k - 1$, and there are no other zeros.

We can now locate the poles of g . If the middle end were flat, then at the level of the end, say $x_3 = c$, the plane $\{x_3 = c\}$ must intersect S in $2kj$ divergent curves, $j \geq 1$. Using this, we can easily show that g has a pole of order $kj + 1$. By (4.1) we can conclude that $j = 1$ and that there are no other poles.

On the other hand, if the middle end is of catenoid-type, the Gauss map has a simple pole there, leaving k more to be found at finite points of S . The planes of symmetry divide S into $2k$ symmetric pieces. If a pole of the Gauss map is found in the interior of one of these regions, there must be at least $2k$ of them. This is a clear contradiction. The sole remaining possibility is to have the poles occurring in the symmetry planes, one on every other of the $2k$ halfplanes.

We may now make a preliminary sketch of the portion of S in a wedge.

The picture suggests that this piece of surface is simply connected. Clearly this surface piece cannot have positive genus, else S would have genus in excess of $2k$, a contradiction.

To show it is simply connected, we will compute its Euler characteristic. By (4.1), its total curvature is $-4\pi(k + 1)/2k$. By taking a sufficiently large piece of the surface we see that each end contributes $\pi + \pi/k$ to the total curvature of the boundary. The first term comes from two right angles at vertices, while the second term comes from the fact that the surface is asymptotic to a catenoid (or plane) and we are seeing a π/k -arc of a circle, which has essentially no normal curvature. All the other boundaries are geodesics. Adding in a

Figure 4.0.1 Successive wedges of S .

contribution of $\pi - \pi/k$ at the single axis point, we have from the Gauss-Bonnet formula

$$\begin{aligned}
 2\pi\chi &= \int K dA + \int k_g + \sum_1^4 \text{vertex angles} \\
 &= \frac{-4\pi(k+1)}{2k} + 0 + 3(\pi + \frac{\pi}{k}) + (\pi - \frac{\pi}{k}) \\
 &= 2\pi.
 \end{aligned}$$

Thus the Euler characteristic of the piece of S in the wedge is 1 and it is simply-connected. We will now show that the sketch is qualitatively correct. There is but one component of the boundary and if we start at the top catenoid end, (considered now in the compactification as a vertex), neither edge emanating from this vertex can go to the bottom catenoid end. This is because the surface is embedded and the top and bottom ends have the same orientation. Hence one edge runs from the top end to the middle end, while the other runs to the axis point. After a Euclidean motion we may assume that if the middle end is a catenoid end, then its logarithmic growth is positive, i.e. it goes up. Thus the two edges emanating from the top catenoid end must look qualitatively like Figure 4.0.1.

We may choose the wedge on the left in this figure. It is clear that there are no other ways to complete the sketch then as in Figure 4.0.1.

4.1 Hidden conformal symmetries

We wish to determine the underlying conformal structure of the surfaces described above. In the previous section, we established that the compactification of the example of genus $k - 1$, $k \geq 2$, assumed to have k vertical planes of symmetry, can be decomposed into $2k$ geodesic 4-gons, all of whose vertex angles equal π/k . All the 4-gons have the same vertices. The Riemann surface must be chosen to have conformal involutions corresponding to the

Figure 4.1 Euclidean rectangles and hyperbolic α -quadrangles.

reflections. If we choose a metric of constant curvature, these conformal involutions are isometries. In the *constant curvature model* our surface is made up of congruent geodesic 4-gons with all angles equal to π/k (rectangles in the case $k = 2$, hyperbolic 4-gons when $k > 2$) and with common vertices. For such 4-gons we need to prove a standard fact from hyperbolic geometry. This will allow us to determine the Riemann surface structure explicitly.

In the case $k = 2$ of genus one, a Euclidean rectangle has two reflective symmetry lines and, conformally, there is precisely a one-parameter family of rectangles. For the case $k > 2$ we will need the same information from hyperbolic geometry about α -quadrangles. By an α -quadrangle we mean a quadrilateral in the hyperbolic plane, all of whose interior angles are equal to α .

Proposition 4.1 *For every $\alpha \in (0, \frac{\pi}{2})$ there exists a 1-parameter family of α -quadrangles. Every α -quadrangle has all the symmetries of a Euclidean rectangle and, for each α , there is a unique α -square, i.e. an α -quadrangle with all the symmetries of a square. (In particular its sides all have the same length.)*

Proposition 4.1 implies that the underlying Riemann surfaces of the minimal surfaces of interest to us have conformal symmetries that are not given by isometries of the minimal surfaces. This is important enough to prove.

Proof of Proposition 4.1. In the Euclidean plane we can parametrize the conformal family of rectangles by the angle $\gamma \in (0, \frac{\pi}{2})$ between the diagonal and an edge. For a given γ , take a triangle with angles $\gamma, \frac{\pi}{2}$ and $\frac{\pi}{2} - \gamma$, and rotate it about the midpoint of the edge opposite the right angle. The two triangles together give the rectangle associated to γ . In the hyperbolic plane: For any angles α, β, γ , the condition $\alpha + \beta + \gamma < \pi$ is necessary and sufficient for the existence of a triangle with these three angles, and, such a triangle is unique up to hyperbolic isometries. Given $\alpha \in (0, \frac{\pi}{2})$, for any $\gamma \in (0, \alpha)$, let $\beta := \alpha - \gamma$ and consider the α, β, γ - triangle. Rotate this triangle about the midpoint of the edge opposite α ; the two triangles together form an α -quadrangle. For each fixed α , different

values of $\gamma \in (0, \alpha)$ give conformally different α -quadrangles because the α, β, γ -triangles are different.

Next, we wish to show that any α -quadrangle has the symmetries of a rectangle. Take any edge, e_1 , of the α -quadrangle, and let σ be the geodesic that is the perpendicular bisector of e_1 . Reflection in σ maps e_1 into itself and interchanges the geodesics that extend the edges, e_0, e_2 , adjacent to e_1 . If the edge e_3 opposite e_1 is symmetric with respect to σ , we are done. If not, e_3 and its σ -reflection \tilde{e}_3 intersect σ and meet the geodesic extensions of e_0 and e_2 at an angle equal to α . This produces two triangles with a common vertex at $e_3 \cap \tilde{e}_3 \in \sigma$ and the same angle $\phi > 0$ there, whose other two angles are α and $\pi - \alpha$, a contradiction since the angle sum is $\pi + \phi > \pi$. Hence $\tilde{e}_3 = e_3$ and the α -quadrangle has the required symmetry.

We wish to show that every α -quadrangle lies in the family constructed above. (Note that our proof that an α -quadrangle has reflectional symmetries did *not* assume that the quadrangle was in this family.) Given an α -quadrangle, the two symmetry geodesics cross at a point, which we will refer to as the *center*. Observe that 180° rotation about the center is a symmetry of the α -quadrangle; it is the composition of the two reflective symmetries. The geodesic join from any vertex to the center extends by this rotation to a diagonal of the α -quadrangle with the center as its midpoint. The diagonal divides the α -quadrangle into two $\alpha, \gamma, \alpha - \gamma$ -triangles for some $\gamma \in (0, \alpha)$. Hence any α -quadrilateral lies in the family we constructed.

If our α -quadrangle was constructed as above with $\gamma = \alpha/2$, note that the diagonal produced from the edge of the $\alpha, \alpha/2, \alpha/2$ triangle (used to construct it) bisects the angles at its endpoints. It follows that reflection in this diagonal is a symmetry of this special α -quadrangle. This is the α -square. \square

In the next section, we will build Riemann surfaces of genus $k - 1$ from $2k$ congruent π/k -quadrangles. Like rectangular tori, they form noncompact 1-parameter families.

4.2 The birdcage model

From the previous two sections we can conclude that the conformal structure of our minimal surface of genus $k - 1$ is given by $8k$ quadrilaterals in hyperbolic (Euclidean) space when $k > 2$ ($k = 2$), with three right angles, one angle of π/k , and edges appropriately identified. In this section, which strictly speaking is not necessary for the next sections, we introduce an embedded surface in 3-space that realizes the conformal symmetries of our minimal surfaces in a geometrically evident manner. It has the same conformal type and will make the assembly of the surface out of quadrilaterals very clear.

Start with k symmetrically placed half-meridians on the unit sphere, $k \geq 2$. Let $S = S_k$ be a tubular neighborhood of these curves in \mathbb{R}^3 . The thickness, $\varepsilon > 0$, of the tubes will correspond to a conformal parameter. The k symmetry planes of the meridians and the

Figure 4.2.0 The birdcage model, $k = 3$, and the fundamental 4-gon.

Figure 4.2.1 The birdcage decomposition The decomposition of the birdcage surface into hyperbolic polygons. Illustration for the case $k = 3$

Upper left: The birdcage with one basic 4-gon darkened. This 4-gon has three right angles and one angle of π/k at an axis point.
Upper right: Four basic 4-gons darkened. Each one has an angle π/k at a different axis point. Taken together, they form a quadrilateral all of whose angles equal π/k .
Lower left: The birdcage decomposed into $2k$ π/k quadrilaterals.
Lower right: The birdcage decomposed into 4 right $2k$ -gons, each centered at an axis point.

equatorial plane slice S into $4k$ geodesic 4-gons. Each 4-gon has two vertices on the polar axis with angle π/k and two vertices on the equator with angle $\pi/2$. For each quadrilateral we have from Gauss-Bonnet that

$$\int K dA = 2\pi - \left[2\left(\frac{\pi}{2}\right) + \frac{2\pi(k-1)}{k} \right],$$

or

$$\int K dA = \pi - \frac{2\pi(k-1)}{k} = -\pi + \frac{2\pi}{k}, \quad (4.2)$$

which implies that the total curvature of S is $-(4k-8)\pi = 2\pi(2-2(k-1))$. This verifies that the genus of S is $k-1$, which is evident from its construction. Note that the reflectional symmetries in the meridian planes generate a k -fold rotational symmetry, and this symmetry fixes the four polar points.

The birdcage surface has the additional isometry of reflection in the equatorial plane. There is another conformal automorphism, which amounts to inversion through the sphere of radius $\sqrt{1-\varepsilon^2}$. These two additional automorphisms restricted to the geodesic 4-gons produce the automorphisms corresponding to the symmetries of a Euclidean rectangle.

Remark 4.1 *The most symmetric of these Riemann surfaces are also the conformal models for one series of Lawson's minimal surfaces in S^3 ([47]). They can be found by extending Plateau solutions, since the symmetry-diagonals of the quadrilaterals, the ones from the π/k -angles, are great-circle arcs of length $\pi/2$ in S^3 .*

The birdcage model serves as a convenient visualization of the decomposition of the Riemann surface into $8k$ 4-gons. We will now work directly with the hyperbolic (or Euclidean when $k=2$) 4-gons having three right angles and one angle of π/k . Note that each of the 4-gons with all angles equal to π/k , decomposes into four 4-gons, each having three right angles and one angle π/k at one of the polar points.

In the next section we construct meromorphic functions on the surface using this decomposition and the Riemann mapping theorem. This will allow us to make the transition from this hyperbolic picture to an algebraic description of our surfaces.

4.3 Meromorphic functions constructed by conformal mappings

Notice that each axis point is the center of a $2k$ -gon, all of whose angles are right angles. These $2k$ -gons are made up of $2k$ of the 4-gons with three right angles, the vertex with angle π/k lying at the polar point. Consider one of these small 4-gons.

We map the quadrilateral into the sector $0 \leq \theta \leq \pi/k$ on the unit disk considered as the closed hemisphere of S^2 by taking the central vertex with angle π/k into 0, its adjacent vertices to 1 and $e^{i\pi/k}$. This determines the image of the opposite vertex, which is a point on the unit circle. We will denote its value by $e^{2i\alpha/k}$, $0 < \alpha < \pi/2$. We will call this map u and extend it to S by Schwarz reflection. Note that at the midpoint of each of the four

Figure 4.3 The conformal mapping used to define u

The illustration here is for $k=3$. The dotted vertex is the branch point of u . Its image is determined by the mapping, and is not assigned in advance.

$2k$ -gons, u has a simple pole or a simple zero, and u maps each of the four $2k$ -gons onto a hemisphere. Thus, u is a degree-2 meromorphic function and we may think of S as a

branched covering of S^2 with $2k$ simple branch points and branch values $e^{i(2\alpha+j\pi)/k}$.

In particular, S is *hyperelliptic*. In the case $k = 2$, u is clearly a degree-2 elliptic function. (In this case it follows directly from the construction of S that S is a rectangular torus.)

We observe that the symmetric construction of u allows us to choose α such that $\pi/4 \leq \alpha < \pi/2$. Given a choice of α between $\pi/4$ and $\pi/2$, simply choose as fundamental domain an adjacent quadrilateral and adjust u by multiplication by $e^{\pi i/k}$.

4.3.1 The function T and its relationship to u

Fix the same quadrilateral on which u is defined as in the previous section. We map this domain onto a quarter-disk, specifying that the vertex opposite the π/k -vertex goes to the origin and the other two $\pi/2$ -vertices are mapped to the unit circle. This determines the mapping up to a rotation in \mathbb{C} , i.e. up to a unitary multiplicative factor. We rotate so that the π/k -vertex is mapped to -1 . We call this mapping T and we label (as in the diagram below) the other two vertices $-e^{-i\beta}$ and $-ie^{-i\beta}$, where the first one is in the upper half-plane.

We will show that, in fact, $\beta = \alpha$, where α is the angle defined by the mapping u . To do this we will construct from T and u , the same conformal map to the lower semi-disk. First of all $(T)^2$ maps our quadrilateral to a semi-disk, which we rotate by $-e^{2i\beta}$ so that $-e^{-i\beta} \rightarrow -1$, $-ie^{-i\beta} \rightarrow +1$, of course $0 \rightarrow 0$, and $-1 \rightarrow -e^{2i\beta}$. In particular, $-e^{2i\beta}(T)^2$ takes the π/k -vertex to $-e^{+2i\beta}$. Turning now to u , note that u^k maps the quadrangle to the upper semi-disk. Composition of u^k with the linear fractional transformation $\nu(\zeta) = \frac{\zeta - e^{2i\alpha}}{1 - \zeta e^{2i\alpha}}$ not only takes the upper semi-disk to the lower semi-disk, but also has the property that

Figure 4.3.1

$\nu(u^k) = -e^{2i\beta}(T)^2$ on the three vertices opposite the π/k -vertex. Thus $\nu(u^k) = -e^{2i\beta}(T)^2$ everywhere, in particular at the π/k -vertex itself; i.e.

$$-e^{+2i\alpha} = \nu(0) = -e^{2i\beta}(-1)^2 = -e^{2i\beta}.$$

Since α and β were both chosen in the range $(0, \frac{\pi}{2})$, $\alpha = \beta$ and

$$\frac{u^k - e^{2i\alpha}}{1 - u^k e^{2i\alpha}} = -e^{2i\alpha}(T)^2.$$

The above equation actually should be considered as the relation between the coordinate maps T and u , and is in fact an equation for our surface.

4.4 The function z and the equation for the Riemann surface in terms of z and u

We prefer to modify T by a Möbius transformation so that: its zeros and poles are more closely related to what we will need in the Weierstrass representation; and it is related to u by an equation with real coefficients. Accordingly let $M(\zeta) = \frac{i(\zeta+1)}{1-\zeta}$ and define

$$z = M \circ T.$$

Note that $T = \frac{z-i}{z+i}$ and that M takes -1 to 0 , 0 to i and 1 to ∞ . Since T takes on the values $-1, 0, -ie^{\pm i\alpha}$ at the vertices of the fundamental 4-gon, we know that z takes on the values 0 and i at two of the vertices. A calculation shows that z takes on the value $\cot(\frac{\pi}{4} + \frac{\alpha}{2})$ at the vertex where T takes on the value $-ie^{-i\alpha}$.¹

¹Let $M(\zeta) = \frac{i(\zeta+1)}{1-\zeta}$. Since $M(e^{i\theta}) = i \cdot \frac{e^{i\theta}+1}{1-e^{i\theta}} = -\frac{e^{i\theta/2}+e^{-i\theta/2}}{(e^{i\theta/2}-e^{-i\theta/2})/i} = -\frac{\cos \theta/2}{\sin \theta/2} = -\cot \theta/2$. It follows that

$$M(-e^{-i\alpha}) = M(e^{i(\pi-\alpha)}) = -\cot(\frac{\pi}{2} - \frac{\alpha}{2}) = -\tan \frac{\alpha}{2},$$

We note that z extends to the $2k$ -gon with a zero of order k at the midpoint, and covers the hemisphere $k/2$ times. Extending to the rest of S by Schwarz reflection produces a function of degree k whose zeros and poles (all of order k) occur at the special polar points. We chose the branch values of u in order to relate u to z in a particularly simple way. We actually have this relationship already because we know that

$$\frac{u^{-k} - e^{2i\alpha}}{1 - u^k e^{2i\alpha}} = -e^{2i\alpha} T^2 = -e^{2i\alpha} \left(\frac{z - i}{z + i} \right)^2.$$

Manipulation of this expression will yield (4.3). We prefer to derive (4.3) by matching the zeros and poles of z and u . Consider $Z := -z + z^{-1}$. From the identity

$$\cot \alpha - \tan \alpha = 2 \cot 2\alpha$$

and the values of z and u given in the previous diagrams, we can easily determine the values of the functions $z^{-1} - z$, and u^{-k} at vertices. Notice that the poles and zeros of $Z + \cot \alpha - \tan \alpha$ match those of u^{-k} and $(\cot \alpha + \tan \alpha)u^{-k}$ equals Z at the points where $u^{-k} = \pm 1$. Hence

$$z^{-1} - z + \cot \alpha - \tan \alpha = (\cot \alpha + \tan \alpha)u^{-k}.$$

This produces an equation for the Riemann surface S , which we write in two equivalent ways:

$$\begin{aligned} z^{-1} - z &= (x + x^{-1})u^{-k} - (x - x^{-1}) \\ z^{-1}(z - x)(z + x^{-1}) &= -(x + x^{-1})u^{-k} \quad \text{where } x = \cot \alpha \end{aligned} \tag{4.3}$$

Here, x may be considered a conformal parameter for our family of Riemann surfaces, $1 \leq x < \infty$.

Differentiate this equation to get

$$-(z + z^{-1}) \frac{dz}{z} = k(x + x^{-1})u^{-k} \frac{du}{u}$$

We now observe that branch points of u occur at points where $z = \pm i$, as we know by construction. By construction or by calculation from equation (4.3) we can conclude that the branch values of u are of the form $e^{\pm 2i\alpha/k} \omega$, where ω is a k^{th} root of unity. At points where $z \neq 0, \infty$, poles of u are branch points of z , with branch values $x = \cot \alpha$ or $-x^{-1} = -\tan \alpha$.

Remark 4.2 *In Section 3, we wrote the above equations in a different and, for historical reasons, more familiar form. Multiply both sides of (4.3) by z^k and define $w := z/u$. Then we may write (4.3) as*

$$\left(\frac{z}{u} \right)^k = w^k = -X^{-1} z^{k-1} (z - x)(z + x^{-1}) \quad X = (x + x^{-1}), \tag{4.4}$$

$$M(e^{i(-\frac{\pi}{2} - \alpha)}) = -\cot(-\frac{\pi}{4} - \frac{\alpha}{2}) = \cot(\frac{\pi}{4} + \frac{\alpha}{2})$$

Figure 4.4 The functions z and u .

Figure 4.4.1

When $k = 2$, the Riemann surface is a rectangular torus. The fundamental domain constitutes $1/16th$ of the surface. Values of u are indicated on the surface. The mapping is specified by fixing the values $-0, 1$, and i at three vertices. It is branched at the fourth vertex; $-$ its value is determined by the mapping, which is in turn determined by the proportions of the rectangle.

and $x, 1 < x < \infty$, parametrizes our family of Riemann surfaces. In case $k = 2$, we have

$$w^2 = -X^{-1}z(z - x)(z + x^{-1}), \quad x \geq 1 \quad (4.5)$$

and we recognize the equation of a rectangular torus. The function z is thus seen to be a geometrically normalized form of the Weierstrass \wp -function: $z = c_1(\wp - c_2)$, where c_2 is such that z has a double zero at the center of one of the 4 gons.

We also note that on the Riemann surfaces (4.4) one can define data for Chen-Gackstatter-type surfaces (see Section 2.2) with higher symmetry using

$$g = \rho \cdot \left(\frac{w}{z}\right)^k = \rho \cdot u^{-k}$$

$$dh = i(z + z^{-1} - 2)\frac{dz}{z},$$

(See Karcher [43].) This construction has been extended by Thayer [69].

4.5 The Weierstrass data

We now determine the Weierstrass data for our minimal surface of genus $k - 1, k \geq 2$, with three vertical catenoid ends and k symmetrically placed vertical planes of reflective symmetry. We know that the underlying Riemann surface will be S , as presented in equation (4.3) or (4.4). Note that we have a 1-parameter family depending on $\pi/4 \leq \alpha < \pi/2$ (or equivalently $x^{-1} = \tan \alpha \geq 1$).

In terms of our decomposition of S , we will look at the region between two adjacent symmetry halfplanes. This corresponds to four basic quadrilaterals joined together to form a π/k -quadrangle with vertices consisting of the polar points of the birdcage where the

vertex angle is π/k . The top and bottom ends are located at points where $u = \infty$. The middle end is placed where $z = \infty$, which is where $u = 0$.

We will specify the Gauss map here by noticing that the values of g must be 0 or ∞ at the ends. (See Figure 4.5.1 and the table in Figure 4.5.0.) Since there is no branching of the Gauss map at a catenoid end these are simple zeros and poles. By contrast, at the saddle we must have at least k curves of intersection with the horizontal tangent plane, which implies that the branching of g at the saddle has order at least $k - 1$. Orient the surface so that $g = 0$ at the saddle. We know that the degree of g must be equal to $k + 1$, and since $g = 0$ at two other catenoid ends, we can conclude that g has a zero of order $k - 1$ at the saddle. Referring to Figure 4.5.0, we note that

$$g_1 := \frac{u^{-1}z}{mz + 1} \quad (4.6)$$

$(z, w) =$	$(0, 0)$	$(x, 0)$	$(-x^{-1}, 0)$	∞, ∞	$z = -m^{-1}$
	Saddle	Bottom End	Top End	Middle End	Finite Vertical Points
g	0^{k-1}	0	0	∞	∞
u^{-1}	∞	0	0	∞	—
z	0^k	x	$-x^{-1}$	∞^k	$-m^{-1}$
$mz + 1$	—	—	—	∞^k	0
dh	0^{k-1}	∞	∞	∞	0
u^k	0^k	∞^k	∞^k	0^k	—
dz/z	∞	0^{k-1}	0^{k-1}	∞	—
du/u	∞	∞	∞	∞	—
$\frac{mz+1}{z+z^{-1}}$	0^k	—	—	—	0

Figure 4.5.0

has the correct zero/pole structure at the saddle and the ends, while having a simple pole at an edge-point in between. Here, m is a free parameter, placing the finite vertical normal at the points where $z = -m^{-1}$. Placement of the vertical normal as it is in the picture requires us to choose $-\infty < (-m^{-1}) < -1/x$, or

$$0 < m < x = \cot \alpha. \quad (4.7)$$

Otherwise, the finite vertical normal would be positioned elsewhere. Fixing m determines the Gauss map up to a real multiplicative factor. (Multiplying by a unitary complex number produces a rotation of the surface.)

$$g = \rho g_1 \quad \rho > 0. \quad (4.8)$$

Figure 4.5.1 The portion of the surface in a wedge and its corresponding π/k -gon.

We are now in a position to determine the complex differential dh of the height function. It must have a simple pole at each catenoid end, a simple zero at the (simple) finite pole of g and a zero of order $k - 1$ at the saddle. Direct calculation of the zeros and poles of $\frac{dz}{z}$, u^k and $mz + 1$ is straightforward. (See Figure 4.5.1.) We may conclude that

$$dh = cu^k(mz + 1)\frac{dz}{z}, \quad (4.9)$$

where c is a nonzero complex multiplicative constant. We wish to show that this constant must be chosen to be real. First, of all we note that the curves on the boundary of this fundamental quadrilateral correspond to planar geodesics (in vertical planes) along which the Gauss map must take values in a meridian circle. Thus $\frac{dg}{g}$ is real on the tangent vectors of these curves. On the other hand, the quadratic differential associated with the second fundamental form of a minimal surface is given as in (2.12) by

$$\langle S(V), V \rangle = \operatorname{Re} \left(\frac{dg(V)}{g} \cdot dh(V) \right). \quad (4.10)$$

Since these planar geodesics are also principal curves, we must have $\frac{dg}{g}(V) \cdot dh(V)$ real along these edges. (See 2.14.) But this implies that dh is real along these edges. From our definitions of u and z , it follows that z and du/u are real along these curves. Hence from (4.10), it follows that c is real. We rescale so that this constant is equal to one. Using (4.3) we may write:

$$dh = u^k(mz + 1)\frac{dz}{z} = \frac{-(x + x^{-1})(mz + 1)dz}{(z - x)(z + x^{-1})}. \quad (4.11)$$

We will use both expressions for dh .

Summary

We summarize what we have achieved.

The Riemann surface and Weierstrass data are as in (4.4), (4.6)–(4.8), (4.11):

$$(x + x^{-1})u^{-k} = z^{-1} - z + (x - x^{-1}) = -z^{-1}(z - x)(z + x^{-1}),$$

$$x = \cot \alpha, \pi/4 \leq \alpha < \pi/2, \text{ minus the points } (z, u) = (x, \infty), (-x^{-1}, \infty)(\infty, 0).$$

The end at $(\infty, 0)$ is the middle end.

$$g = \rho g_1 = \rho \frac{u^{-1}z}{mz + 1} \quad \rho > 0, \quad 0 \leq m < x; \quad (4.12)$$

$$dh = u^k(mz + 1)\frac{dz}{z} = \frac{-(x + x^{-1})(mz + 1)dz}{(z - x)(z + x^{-1})}$$

This data will produce in the Weierstrass Representation

$$X(p) = (x_1, x_2, x_3)(p) = \operatorname{Re} \int_{p_0}^p \Phi, \\ \Phi = \left(\frac{1}{2}(g^{-1} - g), \frac{i}{2}(g^{-1} + g), 1 \right) dh,$$

a (possibly multivalued) minimal immersion into \mathbb{R}^3 . From the development in Sections 4.1 to 4.5, it is clear that if the surfaces $M_{k,x}$ of Theorem 3.3 (also described at the beginning of Section 4) exist, they must have a Weierstrass Representation in the form above, for appropriate choices of constants k, x, m and ρ . All Weierstrass data of the above form produces the symmetry of the surfaces that we require. To see this, first note that the conformal automorphisms

$$\begin{aligned} f(z, u) &:= (\bar{z}, \bar{u}) \\ \tau(z, u) &:= (z, \varepsilon u) \end{aligned} \quad \varepsilon = e^{-\frac{2\pi i}{k}}$$

fix $p_0 := (0, 0)$ and the ends (x, ∞) , $(-x^{-1}, \infty)$ and $(\infty, 0)$. We choose to integrate from this p_0 in the expression for $X(p)$, and observe that we may write

$$\begin{aligned} x_1 - ix_2(p) &= \int_{p_0}^p g^{-1} dh - \overline{\int_{p_0}^p g dh} \\ x_3(p) &= \operatorname{Re} \int_{p_0}^p dh. \end{aligned}$$

Since $f^*(dh) = \overline{dh}$ and $\tau^*(dh) = dh$, it is evident that $x_3(\tau(p)) = x_3(f(p)) = x_3(p)$. Because $f^*g = \bar{g}$ and $\tau^*g = \varepsilon g$, it follows that

$$\begin{aligned} x_1 - ix_2(f(p)) &= \overline{x_1 - ix_2(p)} = x_1 + ix_2(p) \\ x_1 - ix_2(\tau^j(p)) &= \bar{\varepsilon}^j (x_1 - ix_2(p)) = 1 \leq j < k. \end{aligned}$$

Thus f induces reflection in the (x_1, x_3) -plane, while τ^j induces rotation by $2\pi j/k$ about the vertical axis. (We could have used the Schwarz Reflection Principle to prove that f induces a reflection.)

4.6 The logarithmic growth rates

We proceed to determine the logarithmic growth rates of the catenoid ends. From Proposition 2.5*vi*), we know that this growth rate is equal to minus the residue of dh at the end. At the bottom end, where $z = x$, and at the top end, where $z = -x^{-1}$, dz is regular, while at the middle end, where $z = \infty$, dz/z has a simple pole with residue -1 . From (4.12) we can read off the residue of dh at $z = x$ and $z = -x^{-1}$:

$$\begin{aligned} \operatorname{Residue}(dh)_{z=x} &= \frac{-(x + x^{-1})(mx + 1)}{x + x^{-1}} = -(mx + 1) \\ \operatorname{Residue}(dh)_{z=-x^{-1}} &= \frac{-(x + x^{-1})(-mx^{-1} + 1)}{(x^{-1} - x)} = -\frac{m}{x} + 1. \end{aligned} \tag{4.13}$$

Figure 4.6 Logarithmic growth rates at the ends

Because the residues must sum to zero (and dh has poles only at the three ends) we get for the middle end

$$\text{Residue}(dh)_{z=\infty} = m(x + x^{-1}).$$

We could also compute this directly by rewriting the expression for dh in (4.12) in terms of z and expressing it in terms of z^{-1} .

$$dh = \frac{(x - x^{-1})(m + z^{-1})}{z^{-2} + (x - x^{-1})z^{-1} - 1} \frac{dz^{-1}}{z^{-1}}.$$

The logarithmic growth rates are the negatives of these residues: With $m \in [0, x)$ arbitrary, we have $-(mx + 1) < 0 < m(x + x^{-1})$. However, the “middle” end can grow faster than the “top” end, in which case the surface would not be embedded. To avoid this we have the growth rate condition

$$m(x + x^{-1}) < -mx^{-1} + 1. \quad (4.14)$$

4.7 The period and embeddedness problems for the surfaces $M_{k,x}$

We must show that the surfaces $M_{k,x}$ in Theorem 3.3 are in fact well defined by the mapping $X: \Sigma \rightarrow \mathbb{R}^3$ in the Weierstrass representation, (2.7) and (2.8) on the Riemann surface Σ given by (4.12). That is, according to (2.9), we must show that

$$\text{Period}_\alpha(\Phi) = \text{Re} \oint_\gamma \Phi = \text{Re} \oint_\gamma (\phi_1, \phi_2, \phi_3) = \vec{0} \quad (4.15)$$

for all closed curves γ on the punctured Riemann surface. These formulae are restated in the summary at the end of Section 4.5, where we proved that the immersions must have the

required Euclidean symmetry. The built-in symmetry of the mapping is crucial. Consider a fundamental quadrilateral as in Figure 4.7. We label the saddle by S and the top, middle and bottom ends by T, M and B , respectively. Opposite sides of the quadrilateral are mapped into parallel vertical planes by $X = Re \int \Phi$. We require that these planes coincide. After a rotation, we can assume that the curve $X(\widehat{ST})$ lies in the $x_2 = 0$ plane. For $X(\widehat{BM})$ to lie in the same plane, we need only check that

$$Re \oint_{\gamma_1} \phi_2 = 0, \quad (4.16)$$

where γ is the symmetry line of the quadrilateral between edges ST and BM .

The curves $X(\widehat{SB})$ and $X(\widehat{TM})$ lie in planes parallel to the plane $x_2 \cos \pi/k = x_1 \sin \pi/k$. They lie in the same plane if and only if

$$Re \oint_{\gamma_2} (\phi_2 \cos \pi/k - \phi_1 \sin \pi/k) = 0. \quad (4.17)$$

Notice that the symmetry forces $Re \int_{\alpha} \phi_3 = 0$ for every closed curve α on the punctured Riemann surface; we get no conditions from the third differential.

There is no geometric reason why the second period condition (4.17) should be more complicated than the first. In fact it is sufficient to look in detail at the first condition (4.16) and, at the end, handle the second condition by a parameter substitution. To see this, consider the following expressions for ϕ_1 and ϕ_2 .

$$\begin{aligned} 2\rho\phi_1 &= (g_1^{-1} - \rho^2 g_1)dh = (u(\frac{1+mz}{z})^2 - \rho^2 u^{-1})u^k dz \\ 2\rho\phi_2 &= i(g_1^{-1} + \rho^2 g_1)dh = i(u(\frac{1+mz}{z})^2 + \rho^2 u^{-1})u^k dz. \end{aligned} \quad (4.18)$$

If we substitute, in the expression for ϕ_2 , $\tilde{u} = e^{i\pi/k}u$, $\tilde{z} = -z$ and $\tilde{m} = -m$ and define $\tilde{\phi}_2 = \phi_2(\tilde{u}, \tilde{z}, \tilde{m}, \rho)$, we have $\tilde{\phi}_2 = \phi_2 \cos \pi/k - \phi_1 \sin \pi/k$. Thus the second period condition (4.17) can be written in the form

$$Re \oint_{\gamma_2} \tilde{\phi}_2 = 0. \quad (4.17')$$

Now note that \tilde{u} and \tilde{z} satisfy the same Riemann surface equation as u and z , provided we substitute $\tilde{x} = 1/x$ for x :

$$\begin{aligned} (\tilde{x} + \tilde{x}^{-1})\tilde{u}^{-k} &= (x + \frac{1}{x})(-1)u^{-k} = -(z^{-1} - z) - (x - x^{-1}) \\ &= (\tilde{z}^{-1} - \tilde{z}) + (\tilde{x} - \tilde{x}^{-1}). \end{aligned}$$

Notice that $\tilde{x} = x^{-1} = \tan \alpha = \cot(\tilde{\alpha})$, where $\tilde{\alpha} = \pi/2 - \alpha$. Since we will explicitly express (4.16) in terms of m and α , we will be able to express (4.17) by the substitution $m \rightarrow \tilde{m} = -m$ and $\alpha \rightarrow \tilde{\alpha} = \pi/2 - \alpha$. Using (4.18) we may express (4.16) and (4.17) in

Figure 4.7 The period problem

terms of the integrals

$$Q_j = Q_j(m, \alpha) := \int_{\gamma_j} \left(\frac{1 + mz}{z} \right)^2 u^{k+1} dz$$

$$C_j = C_j(\alpha) := \int_{\gamma_j} u^{k-1} dz,$$

$j = 1, 2$; namely

$$Q_1(m, \alpha) = \rho^2 C_1(\alpha)$$

$$Q_2(\tilde{m}, \tilde{\alpha}) = \rho^2 C_2(\tilde{\alpha}) \quad \tilde{m} = -m \quad \tilde{\alpha} = \frac{\pi}{2} - \alpha. \quad (4.19)$$

For each α , we want to find (ρ, m) satisfying these conditions, which express the period problem. Note that Q_j is quadratic in m , while, C_j depends on α only. Each condition in (4.19) determines $\rho^2 = \rho^2(m, \alpha)$, and it is easy to check that $\rho^2 > 0$. The two conditions agree if and only if

$$\frac{Q_1(m, \alpha)}{Q_2(\tilde{m}, \tilde{\alpha})} = \frac{C_1(\alpha)}{C_2(\tilde{\alpha})} \quad (\text{Compatibility condition for } m = m(\alpha)). \quad (4.20)$$

Our strategy is to reduce this condition to a linear equation in $(m - m^{-1})$, whose obvious solvability shows that, for each α , we can find $m(\alpha), \rho(\alpha)$ solving the period problem (4.19). We then need to show that the logarithmic growth rates are correctly ordered. That is, the growth rate at the middle end (the one corresponding to the puncture $(z, u) = (\infty, 0)$) lies between the growth rate at the other two ends. From (4.13) and (4.14), this is equivalent to

$$2 \tan \alpha + \cot \alpha = 2x^{-1} + x < m^{-1} \quad (\text{Growth rate condition for } m = m(\alpha)). \quad (4.21)$$

Proposition 4.2 *Fix $k \geq 2$. For every $\pi/4 \leq \alpha < \pi/2$, there exists a unique non-negative $m(\alpha) < \cot \alpha$ that satisfies the compatibility condition (4.20). Moreover $m(\alpha)$ satisfies the growth condition (4.21) and is a continuous function of α .*

We will give a complete proof of this Proposition in the case $k > 2$. For the torus case, $k = 2$, the arguments are a bit more difficult. (See [30].) As indicated above, existence of the surfaces $M_{k,x}$ in Theorem 3.3 follows from Proposition 4.2. When $\alpha = \frac{\pi}{4}$, $m(\alpha) = m(\frac{\pi}{4}) = 0$, and the middle end is flat; these are the surfaces $M_{k,1}$ of Theorem 3.3, which correspond to the surfaces M_k of Theorem 3.2. These surfaces M_k are embedded. Thus each $M_{k,x}$ lies in a continuous family of examples whose first member is embedded. The embeddedness of the surfaces in the $M_{k,x}$ family is a consequence of the growth condition (4.21) and the following result from [30].

Proposition 4.3 *Suppose $h_t: N \rightarrow \mathbb{R}^3$, $0 \leq t \leq A$ is a continuous one-parameter family of complete minimal immersions of finite total curvature. Suppose that for all t , the ends of $h_t(N) = N_t$ are all vertical, and that h_0 is an embedding but h_A is not. Let $T = \sup\{t | h_t \text{ is an embedding}\}$. Then h_T is an embedding if $T < A$ and if one orders the ends by height, there are at least two ends with the same logarithmic growth.*

4.8 The details of the solution of the period problem, I. Simplification of the integrals.

Recall from (4.3) (or (4.12)) that the Riemann surface equation is

$$z^{-1} - z = (x + x^{-1})u^{-k} - (x - x^{-1}). \quad (4.22)$$

Differentiation gives us a relation between the differentials dz and du :

$$(z^{-1} + z) \frac{dz}{z} = k(x + x^{-1})u^{-k} \frac{du}{u}. \quad (4.23)$$

Two differential forms have the same periods, or are homologous, $w \sim \eta$, if they differ by the differential of a function: $w = \eta + df$. For example:

$$g_1 dh = u^{k-1} dz \sim -(k-1)zu^{k-2} du.$$

This gives us explicit integral expressions for C_1 and C_2 defined in the previous section.

Lemma 4.1

$$C_1 = Re \int_{\gamma_1} i g_1 dh = \frac{2}{k}(k-1)(x + x^{-1}) \int_0^\alpha \sqrt{\cos^2 \phi - \cos^2 \alpha} \cos((1 - \frac{2}{k})\phi) d\phi$$

$$C_2 = \frac{2}{k}(k-1)(x + x^{-1}) \int_0^{\tilde{\alpha}} \sqrt{\cos^2 \phi - \cos^2 \tilde{\alpha}} \cos((1 - \frac{2}{k})\phi) d\phi.$$

Proof. The path γ_1 is symmetric with respect to a branch point of u ; therefore u has the same values at opposite points and du changes sign at the branch point. Moreover z has the value i at this point and is symmetric under 180° rotation, i.e. its values at opposite points are $z, -1/z$. Hence

$$\int_{\gamma_1} 2g_1 dh = -(k-1) \int_{\gamma_1} (z + z^{-1})u^{k-1} \frac{du}{u}.$$

We can eliminate $z + z^{-1}$ as follows:

$$\begin{aligned} \left(\frac{z + 1/z}{x + 1/x} \right)^2 &= \left(\frac{1/z - z}{x + 1/x} \right)^2 + \frac{4}{(x + 1/x)^2} \\ &= \left(u^{-k} - \frac{x - 1/x}{x + 1/x} \right)^2 + \frac{4}{(x + x^{-1})^2}. \end{aligned}$$

Recall $x = \cot \alpha$. Hence $(x - x^{-1})(x + x^{-1})^{-1} = \cos 2\alpha$; with this

$$\left(\frac{z + 1/z}{x + 1/x} \right)^2 = u^{-k} \cdot (u^k + u^{-k} - 2 \cos 2\alpha). \quad (4.24)$$

Furthermore, along the path γ_1 we can take u as coordinate function on the Riemann surface; since u is unitary along γ_1 we have

$$u = e^{i\phi}, u^k + u^{-k} = 2 \cos \phi, \quad 0 \leq \phi \leq \frac{2\alpha}{k}$$

$$du/u = id\phi.$$

We insert this and (4.24) in the last integral and get

$$Re \int_{\gamma_1} i g_1 dh = (k-1)(x+x^{-1}) \int_0^{2\alpha/k} \sqrt{2 \cos k\phi - 2 \cos 2\alpha} \cos((\frac{k}{2}-1)\phi) d\phi.$$

We substitute $t = k\phi$ and use $\cos 2\alpha = 2\cos^2 \alpha - 1$ to derive the expression for C_1 in the lemma. Using the symmetric treatment discussed above, we get C_2 if we replace α by $\tilde{\alpha} := \frac{\pi}{2} - \alpha$. \square

Now we turn to $Q_1(m)$. We have at first

$$\frac{1}{g_1} dh = (m^2 + \frac{1}{z^2} + \frac{2m}{z}) u^{k+1} dz,$$

so that $\int_{\gamma_1} \frac{1}{g_1} dh$ is given in terms of three different period integrals on the Riemann surface. We have to reduce these integrals to the two in the previous lemma and to two others before we can discuss them. First, multiply the differential relation (4.23) by $u^{k+\ell}$ to get

$$u^{k+\ell}(1+z^{-2})dz = k(x+x^{-1})u^\ell \cdot \frac{du}{u} \underset{[\ell \neq 0]}{\sim} 0.$$

In particular,

$$\frac{1}{z^2} u^{k+1} dz \sim -u^{k+1} dz.$$

Hence we achieved a first reduction to two integrals per path:

$$\frac{1}{g_1} dh \sim (m^2 - 1 + \frac{2m}{z}) u^{k+1} dz.$$

We deal with $u^{k+1} dz \sim -(k+1)u^{k+1} z du/u$ as before by using the symmetry along γ_1 :

$$\int_{\gamma_1} 2u^{k+1} dz = - \int_{\gamma_1} (k+1)u^{k+1} (z + \frac{1}{z}) \frac{du}{u}.$$

Again $(z + 1/z)$ is eliminated using (4.24) to get almost the same integral as in the lemma:

$$2Re \int_{\gamma_1} i u^{k+1} dz = \frac{4(k+1)}{k} (x+x^{-1}) \int_0^\alpha \sqrt{\cos^2 \phi - \cos^2 \alpha} \cdot \cos((1 + \frac{2}{k})\phi) d\phi. \quad (4.25)$$

The remaining integral $\int u^{k+1} dz/z$ can be rewritten as a linear combination of the integrals we already have. We multiply the differential relation (4.23) to get

$$(z + \frac{1}{z}) u^{k+1} dz = k(x+x^{-1}) z du$$

and we multiply the surface equation (4.22) to get

$$(\frac{1}{z} - z) u^{k+1} dz = (x+x^{-1}) u dz + (x^{-1} - x) u^{k+1} dz.$$

Before we add these two equations, we use $udz \sim -zdu$. Then

$$\frac{2}{z}u^{k+1}dz \sim (k-1)(x+x^{-1})zdu + (x^{-1}-x)u^{k+1}dz.$$

The second differential on the right has been handled in (4.25). The remaining one is treated with path-symmetry as before:

$$\int_{\gamma_1} 2zdu = \int_{\gamma_1} (z + \frac{1}{z})du,$$

and $(z + 1/z)$ is eliminated with (4.24). We obtain

$$2Re \int_{\gamma} izdu = -\frac{4}{k}(x+x^{-1}) \int_0^{\alpha} \sqrt{\cos^2 \phi - \cos^2 \alpha} \cos((-1+2/k)\phi) d\phi. \quad (4.26)$$

We do not know an a priori reason why we arrived at the same integral as in the lemma, but we now can use this fact to finish the period discussion. First shorten the notation by defining

$$\begin{aligned} E_{\mp}(\gamma) &:= \int_0^{\gamma} \sqrt{\cos^2 \phi - \cos^2 \gamma} \cdot \cos((1 \mp 2/k)\phi) d\phi, \\ c &:= \frac{2}{k}(x+x^{-1}). \end{aligned} \quad (4.27)$$

In these terms the lemma says

$$C_1 = c \cdot (k-1) \cdot E_-(\alpha), \quad C_2 = c \cdot (k-1) \cdot E_-(\tilde{\alpha}).$$

The simplifications concentrated in (4.25), (4.26) deal with all the terms in

$$\frac{1}{g_1}dh \sim (m^2 - 1 + \frac{2m}{z})u^{k+1}dz$$

to give

$$\begin{aligned} Q_1(m) &:= Re \int_{\gamma_1} -i \frac{dh}{g_1} \\ &= c \cdot [(1-m^2)(k+1)E_+(\alpha) + (\cot \alpha - \tan \alpha) \cdot m \cdot (k+1)E_+(\alpha) \\ &\quad + (\cot \alpha + \tan \alpha) \cdot m \cdot (k-1)E_-(\alpha)]. \end{aligned}$$

The substitution $\alpha \rightarrow \tilde{\alpha} := \pi/2 - \alpha$, $m \rightarrow \tilde{m} := -m$ gives

$$\begin{aligned} Q_2(m) &= c \cdot [(1-m^2)(k+1)E_+(\tilde{\alpha}) + (\cot \alpha - \tan \alpha) \cdot m \cdot (k+1)E_+(\tilde{\alpha}) \\ &\quad - (\cot \alpha + \tan \alpha) \cdot m \cdot (k-1)E_-(\tilde{\alpha})]. \end{aligned}$$

The compatibility condition (4.20) is now fractional linear in $M := \frac{1}{m} - m$ and of the form

$$\frac{M + a + b \cdot q}{M + a - b \cdot \tilde{q}} = \frac{q}{\tilde{q}},$$

namely:

$$\frac{(\frac{1}{m} - m) + (\cot \alpha - \tan \alpha) + (\cot \alpha + \tan \alpha) \frac{k-1}{k+1} \cdot \frac{E_-(\alpha)}{E_+(\alpha)}}{(\frac{1}{m} - m) + (\cot \alpha - \tan \alpha) - (\cot \alpha + \tan \alpha) \frac{k-1}{k+1} \cdot \frac{E_-(\alpha)}{E_+(\alpha)}} = \frac{E_-(\alpha)/E_+(\alpha)}{E_-(\tilde{\alpha})/E_+(\tilde{\alpha})}$$

This simplifies to

$$((\frac{1}{m} - m) + (\cot \alpha - \tan \alpha)) \cdot \frac{k+1}{k-1} \cdot \left(\frac{E_+(\tilde{\alpha})}{E_-(\tilde{\alpha})} - \frac{E_+(\alpha)}{E_-(\alpha)} \right) = 2 \cdot (\cot \alpha + \tan \alpha). \quad (4.28)$$

4.9 The details of the solution of the period problem, II. The monotonicity lemma.

We will prove the following

Lemma 4.2 (The Monotonicity Lemma) *Fix $k \geq 2$. The function $\gamma \rightarrow \frac{E_+(\gamma)}{E_-(\gamma)}$ is strictly decreasing on $(0, \frac{\pi}{2})$, with*

$$\lim_{\gamma \rightarrow 0} \frac{E_+(\gamma)}{E_-(\gamma)} = 1, \quad \lim_{\gamma \rightarrow \frac{\pi}{2}} \frac{E_+(\gamma)}{E_-(\gamma)} = \frac{k-1}{k+1}.$$

The integrals $E_{\pm}(\gamma)$ are defined in (4.27).

Before presenting the proof, we will show how the Lemma is used to prove Proposition 4.2. Recall from Section 4.3 that the Riemann surfaces determined by the conformal parameter α , with $\alpha > \pi/4$, are obtained from those with $\alpha < \pi/4$ by exchanging generators. The choice of $\alpha > \pi/4$ was made to be compatible with the location of the point where the normal is vertical (i.e. where $z = -m^{-1}$) on the planar geodesic symmetry curve *between* the end-punctures at the points where $z = -x^{-1} = -\tan \alpha$ and $z = \infty$. To see this, let $\alpha > \pi/4$. Then the Monotonicity Lemma implies

$$\frac{2}{k+1} = \frac{E_+}{E_-}(0) - \frac{E_+}{E_-}\left(\frac{\pi}{2}\right) \geq \frac{E_+(\tilde{\alpha})}{E_-(\tilde{\alpha})} - \frac{E_+(\alpha)}{E_-(\alpha)} > 0.$$

With this, it is clear that (4.28) is *always* solvable for $m^{-1} - m$ and that (for $\alpha > \pi/4$)

$$m^{-1} - m > (k-1)(\cot \alpha + \tan \alpha) + (\tan \alpha - \cot \alpha) > 0 \quad (4.29)$$

$$m^{-1} - \tan \alpha + (\cot \alpha - m) > (k-1)(\cot \alpha + \tan \alpha) > 0.$$

This shows that $M := m^{-1} - m > 0$, that $m = -\frac{M}{2} + \sqrt{1 + \frac{M^2}{4}} < 1/m$, and that $m < \cot \alpha$. Therefore the positive solution $m = m(\alpha)$ of (4.28) (which is our compatibility condition (4.20)), puts the vertical normal on the symmetry line defined by $-\infty < z < -\tan \alpha = -x^{-1}$, that runs between the punctures at $z = -\tan \alpha$ and $z = \infty$. Thus we have established the first part of Proposition 4.2 for any $k \geq 2$. For $k > 2$, the inequality (4.29) has as an immediate consequence the inequality

$$m^{-1} > 2 \tan \alpha + \cot \alpha = 2x^{-1} + x,$$

which is the growth rate condition (4.21). This concludes the proof of Proposition 4.2, when $k > 2$.

Remark 4.3 *The case $k = 2$, the genus=1 case, turned out to be more difficult for us than the higher genus cases. In addition to the Monotonicity Lemma, we proved, for $k = 2$:*

$$\frac{E_+(\tilde{\alpha})}{E_-(\tilde{\alpha})} - \frac{E_+(\alpha)}{E_-(\alpha)} \leq \frac{2}{3} \sin^2 \alpha.$$

If we insert this in the compatibility condition (4.28), and use the identity $\sin^{-2} \alpha = \cot^2 \alpha + 1$, we get, instead of (4.29),

$$\begin{aligned} m^{-1} - m &> \frac{\cot \alpha + \tan \alpha}{\sin^{-2} \alpha} - \cot \alpha + \tan \alpha \\ &= \cot^3 \alpha + \tan \alpha [\cot^2 \alpha + 2] \\ &= \cot^3 \alpha + \cot \alpha + 2 \tan \alpha \\ &> 2 \tan \alpha + \cot \alpha = 2x^{-1} + x, \end{aligned}$$

which implies the growth rate condition (4.21) in the case $k = 2$.

4.9.1 Proof of the Monotonicity Lemma

To establish the claim

$$0 \leq \alpha \leq \gamma \leq \frac{\pi}{2} \Rightarrow \frac{E_+}{E_-}(\alpha) \geq \frac{E_+}{E_-}(\gamma)$$

it is equivalent to prove

$$E_-(\alpha) \cdot (E_-(\gamma) - E_+(\gamma)) - E_-(\gamma) \cdot (E_-(\alpha) - E_+(\alpha)) \geq 0 \quad (4.30)$$

and sufficient to prove

$$E_-(\alpha) \cdot (E_- - E_+)'(\alpha) - E_- '(\alpha) \cdot (E_- - E_+(\alpha)) \geq 0. \quad (4.31)$$

From the definition of these integrals in (4.27) we have

$$E_- '(\alpha) = \sin \alpha \cos \alpha \int_0^\alpha \frac{1}{\sqrt{\cos^2 \phi - \cos^2 \alpha}} \cos\left(\left(1 - \frac{2}{k}\right)\phi\right) d\phi$$

and using $\cos(\alpha - \beta) - \cos(\alpha + \beta) = 2 \sin \alpha \sin \beta$ we have

$$(E_- - E_+)(\alpha) = \int_0^\alpha \sqrt{\cos^2 \phi - \cos^2 \alpha} \cdot 2 \sin \phi \cdot \sin\left(\frac{2}{k}\phi\right) d\phi,$$

$$(E_- - E_+)'(\alpha) = \sin \alpha \cos \alpha \int_0^\alpha \frac{2 \sin \phi \cdot \sin\left(\frac{2}{k}\phi\right)}{\sqrt{\cos^2 \phi - \cos^2 \alpha}} d\phi.$$

Now we define decreasing functions f_1, g_1 and increasing functions f_2, g_2 with choices of constants $c_i > 0$ such that $\int_0^\alpha g_i(\phi) d\phi = 1$:

$$\begin{aligned} 0 \leq f_1(\phi) &:= \sqrt{\cos^2 \phi - \cos^2 \alpha} \\ 0 \leq g_1(\phi) &:= c_1 \cdot \cos\left(\left(1 - \frac{2}{k}\right)\phi\right) \quad (\text{decreasing}) \\ 0 \leq f_2(\phi) &:= 1/\sqrt{\cos^2 \phi - \cos^2 \alpha} \\ 0 \leq g_2(\phi) &:= c_2 \cdot 2 \sin \phi \cdot \sin\left(\frac{2}{k}\phi\right) \quad (\text{increasing}). \end{aligned}$$

With this notation (4.31) is equivalent to

$$\int_0^\alpha f_1 \cdot g_1 d\phi \cdot \int_0^\alpha f_2 \cdot g_2 d\phi \geq \int_0^\alpha f_1 \cdot g_2 d\phi \cdot \int_0^\alpha f_2 \cdot g_1 d\phi$$

and this inequality is implied by

$$\int_0^\alpha f_1 \cdot (g_1 - g_2) d\phi \geq 0, \quad \int_0^\alpha -f_2 \cdot (g_1 - g_2) d\phi \geq 0.$$

Since $g_1 - g_2$ is decreasing, with $\int_0^\alpha (g_1 - g_2) = 0$, there exists a $\phi^* \in (0, \alpha)$ such that

$$\begin{aligned} \phi \leq \phi^* &\Rightarrow (g_1 - g_2)(\phi) \geq 0, \\ f_1(\phi) &\geq f_1(\phi^*) \geq 0 \quad \text{and} \quad -f_2(\phi^*) \leq -f_2(\phi) \leq 0 \\ \phi^* \leq \phi &\Rightarrow (g_1 - g_2)(\phi) \leq 0 \\ 0 \leq f_1(\phi) &\leq f_1(\phi^*) \quad \text{and} \quad -f_2(\phi) \leq -f_2(\phi^*) \leq 0. \end{aligned}$$

Therefore

$$\begin{aligned} \int_0^\alpha f_1(\phi)(g_1 - g_2)(\phi) d\phi &\geq \int_0^\alpha f_1(\phi^*)(g_1 - g_2)(\phi) d\phi = 0 \\ \int_0^\alpha -f_2(\phi)(g_1 - g_2)(\phi) d\phi &\geq \int_0^\alpha -f_2(\phi^*)(g_1 - g_2)(\phi) d\phi = 0. \end{aligned}$$

□

5 The structure of the space of examples

We begin this section by giving a list of the constructions to date of complete embedded minimal surfaces with finite total curvature. None of these examples have been as completely analyzed as those of Theorem 3.3. For some, a complete existence proof independent of computation has yet to be found. For others there are existence proofs that amount to computing, with error estimates, the degree of the period map. All of these examples have the property that they are part of deformation families of surfaces, which begin with all but two of the ends flat and continue through surfaces all of whose ends are catenoidal. Most of the families, in contrast to the surfaces $M_{k,x}$ of Theorem 3.3 eventually contain surfaces that are not embedded because the growth rates of the ends change their order and a lower end overtakes a higher one.

Figure 5.0.1 Four-ended embedded minimal surfaces of genus $2k, k > 0$.
These are the surfaces described in 1).

Figure 5.0.2 Four-ended embedded minimal surfaces.

Right: The surface described in 3) with $k = 1$.
Left: Deformation of the four-ended surface on the left, now with all ends catenoidal. The surface is no longer embedded.

Figure 5.0.3 Five-ended embedded minimal surface of genus 3,
discovered by Boix and Wohlgemuth. See 4).

- 1) Four ended examples with $k \geq 2$ vertical planes of symmetry, one horizontal plane of symmetry, genus $2(k - 1)$, two flat ends and two catenoid ends [73]. (See Figure 5.0.1.)
- 2) A one parameter family deforming the surfaces above through surfaces with four catenoid ends. Eventually these surfaces cease to be embedded. [72, 4, 73, 74].
- 3) The surfaces in 2) with symmetric tunnels through their waist planes [72, 4]. These surfaces have $k \geq 2$ vertical planes of symmetry and genus $3(k - 1)$. The family begins with a surface with two flat ends and moves through surfaces with all four ends catenoidal. Eventually the surfaces are not embedded. (See Figure 5.0.2.)
- 4) Five ended examples. [5]. (See Figure 5.0.3.)

These examples have in common the fact that they were discovered first with the help of insights guided by and guiding computation. The software MESH, developed by James T. Hoffman et. al. ([8, 41]) together with various numerical routines for killing periods (finding simultaneous zeros of several real-valued functions of several variables) has been crucial to this research. This was also the case for the three-ended examples of Theorem 3.3. One

has to produce Weierstrass data by methods analogous to those presented in Section 4.1–4.4. The main obstacle to constructing more complex examples with methods like those in Section 4 is the difficulty in killing the periods. Even numerically, for n large, finding the simultaneous zero of n functions on a domain in \mathbb{C}^n is time-consuming and unstable for functions that themselves are approximated by numerical integration (the periods given by Weierstrass integrals).

The state of knowledge at present is rather poor. It is not known whether these surfaces can be perturbed further to reduce their symmetry. There is at present no general implicit-function-theorem type result for the period mapping. Some important theoretical progress has been made by Wohlgemuth [73, 74]. The existence of certain of these surfaces (the most symmetric ones with flat ends) can be proved by theoretical means involving finding desingularizing limits of unstable solutions to the Plateau Problem. (See, for example [34].) However, where this is done for the surfaces of Theorem 3.2, this procedure has never produced an example not previously discovered by the method outlined above.

5.1 The space of complete embedded minimal surfaces of finite total curvature

Let \mathcal{M} be the space of complete embedded minimal surfaces of finite total curvature, normalized so that all the ends have vertical normals and so that the maximum absolute value of the Gauss curvature is equal to one and occurs at the origin. (Note that the plane is not in \mathcal{M} .)

Definition 5.1 *The subspace $\mathcal{M}_{i,j} \subset \mathcal{M}$, $i \geq 0, j \geq 2$, consists of all surfaces in \mathcal{M} of genus i with j ends. We define*

$$\mathcal{M}_{k,r}^* = \bigcup_{i \leq k, j \leq r} \mathcal{M}_{i,j}.$$

We consider two elements of \mathcal{M} to be the same if they differ by a Euclidean motion. From (2.21) it follows that the total curvature of an example in \mathcal{M}_{ij} is equal to $-4\pi(1 - (i + j))$. This means that the subspace of \mathcal{M} with total curvature at least $-4\pi n$ is a union of a finite number of $\mathcal{M}_{k,r}^*$. In this terminology, we may restate some of Theorem 3.1 and Theorem 3.4 as follows:

- i) $\mathcal{M}_{0,r}^*$ consists of one surface, the catenoid [51];
- ii) $\mathcal{M}_{k,2}^*$ consists of one surface, the catenoid [66];
- iii) $\mathcal{M}_{1,3}$ consists of the surfaces $M_{2,x}$ in Theorem 3.3, which is naturally a half-open interval $[1, \infty)$ [16].

Note from this last item that the spaces \mathcal{M}_{ij} are not necessarily compact. However it can be shown that in \mathcal{M} , which we consider with the compact-open topology, every

divergent sequence of surfaces $M_{k,x}$ has a subsequence that converges to the catenoid [30]. Thus while $\mathcal{M}_{1,3}$ is not compact, $\mathcal{M}_{1,3}^*$ is naturally a closed interval and is compact. (Its end points are Costa's surface and the catenoid.) The following result generalizes this fact.

Theorem 5.1 ([30]) $\mathcal{M}_{k,r}^*$ is compact. Specifically, every sequence of surfaces in $\mathcal{M}_{p,k}^*$ possesses a subsequence that converges smoothly on compact subsets of \mathbb{R}^3 to a surface in $\mathcal{M}_{p,k}^*$.

This theorem is of limited utility because at the present time it is not known, in any general fashion, which of the spaces $\mathcal{M}_{i,j}$ are empty and which are not.

5.2 Some questions and conjectures

One could imagine that the Costa example is constructed in the following heuristic manner. Take a catenoid and intersect it by its waist plane. Remove a tubular neighborhood of the intersection circle and smoothly join up the four boundary circles by a genus one minimal surface. The higher-genus examples $M_k, k \geq 3$ can be thought of as improved approximations.

Theorem 5.2 ([38])

- i) As normalized in Theorem 3.2, the surfaces $M_k, k \geq 2$, possess a subsequence that converges as $k \rightarrow \infty$ to the union of the plane and the catenoid. The convergence is smooth away from the intersection circle.
- ii) Considered as surfaces in \mathcal{M} (i.e., normalized so that the maximum value of K is equal to 1 and occurs at the origin), the sequence $\{M_k\}$ possesses a subsequence that converges to Scherk's singly-periodic surface.

It has been conjectured by Hoffman and Meeks that for a complete embedded nonplanar minimal surface of finite total curvature

$$k \geq r - 2 \tag{5.1}$$

where k is the genus and r is the number of ends. Theorem 3.1, Statements 2 and 3, results of Lopez-Ros and of Schoen, can be interpreted as saying that equality holds in (5.1) when either $k = 0$ or $r = 2$. The examples of Theorems 3.2 and 3.3, as well as all other known examples satisfy this inequality. Inequality 5.1 may be stated as a conjecture

Conjecture 5.1

$$\mathcal{M}_{k,r} \text{ is empty if } k < r - 2.$$

Figure 5.2.0

One of a family of complete minimal surfaces of genus zero with two parallel, embedded catenoid ends and one embedded flat end, at which the Gauss map has an order-two branch point. The surface is not embedded because the flat end is not parallel to the catenoid ends.

The heuristic idea behind the conjecture is that to increase the number of ends (r) by adding a flat end and then looking for a Costa style surface like the surface M_k , which is nearby and embedded, forces the genus (k) to increase. Related to this conjecture is the question of whether the order of the Gauss map at a flat end of a complete embedded minimal surface can be equal to two. According to Remark 2.3iii) the order must be at least two, and in all known *embedded* examples of finite total curvature the order is at least three.

(See Remark 2.8.) However, there are complete embedded periodic minimal surfaces with flat ends where the Gauss map has degree two. The most famous of these is the example of Riemann [62, 63]. (See also [31], [37] and [39].) For *immersed* minimal surfaces, it is possible to have such an end. Consider the Weierstrass data

$$g(z) = ((z - r)(z + r))^{-1}, dh = (z^2 - r^2)(z^2 - 1)^{-2}dz, r \neq 1,$$

on $S^2 - \{\pm 1, \infty\}$. The surface has sufficient symmetry (vertical plane of reflection and one horizontal line) to make the period problem one-dimensional. The period problem is solvable by a residue calculation, which determines the choice of r . The ends at ± 1 are catenoid ends and the end at $g(z) = \infty$, where $g(z)$ has a zero of order 2, is flat. However, while all the ends are separately embedded, they are not parallel; the surface is complete and immersed but not embedded.

Figure 5.2.1 The Horgan Surface

The surface depicted here with genus two and three ends probably does not exist. The periods appear to be zero, but one of them is just very small. Computationally, it appears that this period always has the same sign and is never zero, although it becomes vanishingly small. However, in the limit, the surface degenerates.

Recall from Proposition 2.1) and Remark 2.4*i*) that each embedded catenoid end of a complete embedded minimal surface of finite total curvature is asymptotic to the end of a specific vertical catenoid. There are always at least two such ends, according to Proposition 2.5*v*) and the discussion following Theorem 2.4. Each such end has an axis determined by the term of order $\mathcal{O}(\rho^{-1})$ in (2.21) and made precise in Definition 2.7. Because all of the known examples can be constructed by symmetry methods like those used in Section 4, it follows that all of the axes of the catenoid ends of these examples coincide. One basic question is whether or not there exist examples whose catenoid ends have *different* axes. Such an example could have at most, a symmetry group of order four, generated by a reflection through a plane containing all of the axes (if such a plane exists) and an order two rotation about a horizontal line (if such a line exists). If the axes were not coplanar, only the rotation could exist. Such a rotation would imply that the number of catenoid ends was even.

As observed in Remark 2.8, if there are three catenoid ends then their axes are coplanar and the end whose axis is in the middle has logarithmic growth with sign opposite that of the other two.

It is known that any intrinsic isometry of a complete *embedded* minimal surface of finite total curvature in \mathbb{R}^3 must extend to a symmetry of that surface; that is an isometry of \mathbb{R}^3 that leaves the surface invariant [13, 36]. If the surface is not embedded, this is

not necessarily the case. The Enneper surface and its generalizations $g(z) = z^k, dh = g(z)dz, z \in \mathbb{C}$, have intrinsic isometric rotations $z \rightarrow uz, |u| = 1$, but only k of them extend to symmetries of the surface. (See Remark 2.3ii).

An even more basic problem concerns the question of whether or not there exists a complete embedded minimal surface with no symmetries except the identity. That all the known examples have significant symmetry groups is an artifact of their construction and says nothing about the general situation. It would be extremely surprising if the hypotheses of finite total curvature together with embeddedness implied the existence of a nontrivial symmetry.

A related question concerns the surfaces $M_{k,x}$ in Theorem 3.3. Costa's Theorem 3.1.5 states that for the genus-one examples ($k = 2$)

$$\mathcal{M}_{1,3} = \{M_{2,x} \mid 1 \leq x < \infty\}. \quad (5.2)$$

It would be very interesting to know to what extent this is true for higher genus:

$$\mathcal{M}_{k,3} \stackrel{?}{=} \{M_{k+1,x} \mid 1 \leq x < \infty\}. \quad (5.3)$$

It is not even known if this is true with the added hypothesis that the symmetry group be of order at least $2(k+1)$, as is the case for the surfaces $M_{k+1,x}$. Costa's proof of (5.2) uses the full force of elliptic function theory, and hence is restricted to genus 1. A geometric proof of (5.2) would be extremely useful to have, in order to begin to understand what happens in the higher-genus, three-ended case.

6 Finite total curvature versus finite topology

A surface has finite topology provided it is homeomorphic to a compact Riemann surface from which a finite number of points have been removed. As a consequence of Theorem 2.2, we know that a complete immersed minimal surface whose total curvature is finite must have finite topology. In fact, it is *conformal* to a punctured compact Riemann surface. In this section, we will discuss to what extent finite topology implies finite total curvature for complete properly embedded minimal surfaces.

The helicoid is a complete, embedded, simply-connected (genus zero and one end) minimal surface. Because it is periodic and not flat, its total curvature is infinite. This simple classical example shows that finite topology does not imply finite total curvature. Osserman has posed the question of whether or not the helicoid is the only nonplanar, complete, embedded, minimal surface of genus-zero with one end. According to Theorem 3.1 there is no such surface of finite total curvature of any genus. One could ask, more generally, if there are any complete embedded minimal surfaces of finite topology and one end, besides the plane and the helicoid. For example, does there exist such a surface that is morphologically

the helicoid with a finite number of handles? (We have recently shown that the answer is yes. See Section 6.3.)

An end of a complete surface of finite topology is necessarily annular. That is, it is homeomorphic to a punctured disk. If it is conformal to a punctured disk and the total curvature is infinite, then the Gauss map must have an essential singularity. All known embedded annular ends are asymptotic to the end of a helicoid. No embedded end is known that is conformal to an annulus $r_1 \leq |z| < r_2$.

6.1 Complete, properly-immersed, minimal surfaces with more than one end

Under the assumption that the surface has more than one end, much more is known.

Theorem 6.1 (Hoffman and Meeks [35]) *Let M be a complete properly embedded minimal surface in \mathbb{R}^3 with more than one topological end. Then at most two of the annular ends have infinite total curvature.*

Corollary 6.1 ([35]) *Let M be a complete embedded minimal surface in \mathbb{R}^3 with finite topology. Then all except two of the ends have finite total curvature and are asymptotic to planes or catenoids, all with parallel limit normals.*

This Corollary was strengthened by Fang and Meeks who proved

Theorem 6.2 (Fang and Meeks [21]) *Let M be a properly embedded minimal surface in \mathbb{R}^3 with two annular ends that have infinite total curvature. Then these ends lie in disjoint closed halfspaces and all other annular ends are flat ends parallel to the boundary of the halfspaces.*

Corollary 6.2 *Suppose M is a complete properly embedded minimal surface of finite topology. If M does not have finite total curvature, then there are three possibilities:*

- i) M has a catenoid end and exactly one end with infinite total curvature that lies in a halfspace;*
- ii) M has exactly one end (of infinite total curvature) and this end does not lie in a halfspace;*
- iii) M has two ends of infinite total curvature that lie in disjoint closed halfspaces, and a finite number of flat ends in the slab between these halfspaces. There are no catenoid ends.*

In particular a complete properly embedded minimal surface of finite topology with more than one end can never have an end asymptotic to a helicoid end (because such an end never lies in a halfspace).

6.2 Complete embedded minimal surfaces of finite topology and more than one end

What sorts of infinite total curvature ends can appear on a complete embedded minimal surface with finite topology? Nitsche considered ends that were fibred by embedded Jordan curves in parallel planes. These are called *Nitsche ends* by Meeks and Rosenberg [53]. For concreteness we assume that such an end A lies in the halfspace $\{x_3 \geq 0\}$ with boundary a simple closed curve in $x_3 = 0$, and that $\{x_3 = t\} \cap A$ is a simple closed curve, for all $t \geq 0$. Nitsche proved

Proposition 6.1 ([57]) *A Nitsche end, all of whose level curves are star shaped, is a catenoidal end.*

In particular, it has finite total curvature. A Nitsche end necessarily has the conformal type of a punctured disk. Certainly, we may assume that the end is conformally $\{z | 0 \leq \alpha < |z| \leq 1\}$. The height function $h(z) = x_3(z)$ is harmonic and so is essentially determined by its boundary values. Suppose $\alpha > 0$. Then $h = 0$ on $|z| = 1$ and, $h = \infty$ on $|z| = \alpha$. Since $c_1(\ln |z|)$ is also harmonic, and $c_1 \ln |z|$ lies above h on the boundary of $\alpha < |z| \leq 1$, it follows that

$$h(z) < c_1 \ln |z| \quad \alpha < |z| < 1$$

for any real value of c_1 . This means $h(z) \equiv -\infty$ on the interior of the annulus, which is a contradiction. Hence we can take the disk to be $\{z | 0 < |z| \leq 1\}$, and $h(z) = c_1 \ln |z|$, $c_1 < 0$. Thus dh is determined and using the Weierstrass representation (2.7), (2.8), we can determine the end completely by the knowledge of its Gauss map. For the Nitsche end to have infinite total curvature, g must have an essential singularity. Thus finding a non-catenoidal Nitsche end is equivalent to finding an analytic function on the punctured unit disk with an essential singularity at the origin for which (2.7) (2.8) provide a well-defined embedding. Toubiana and Rosenberg have given examples which are well-defined, but not embedded [64]. We also remark that on \mathbb{C} , $g(z) = \exp(F(z))$, and $dh = dz$, where $F(z)$ is entire, gives a complete, immersed, simply-connected minimal surface whose Gauss map has an essential singularity at infinity.

The importance of understanding Nitsche ends is underscored by the following strong result of Meeks and Rosenberg.

Theorem 6.3 ([53]) *Let M be a complete embedded minimal surface of finite topology and more than one end. Then any end of infinite total curvature is a Nitsche end. In particular M is conformal to a compact Riemann surface punctured in a finite number of points.*

As a result of this theorem, a proof of the Nitsche conjecture would imply that finite topology and finite total curvature are equivalent for complete embedded minimal surfaces with at least two ends.

Figure 6.3 The genus one helicoid, $\mathcal{H}e_1$

6.3 Complete embedded minimal surfaces of finite topology with one end

Very recently, Hoffman, Karcher and Wei [32, 33] have constructed a complete embedded minimal surface of finite topology and infinite total curvature.

Theorem 6.4 *There exists a complete embedded minimal surface, $\mathcal{H}e_1$, of genus-one with one end that is asymptotic to the end of a helicoid. The surface contains two lines, one vertical and corresponding to the axis of the helicoid, the other horizontal crossing the axial line. Rotation about these lines generates the full isometry group of $\mathcal{H}e_1$, which is isomorphic to $Z_2 \oplus Z_2$.*

This is the first properly embedded minimal surface with infinite total curvature, whose quotient by symmetries does not have finite total curvature.

7 Stability and the index of the Gauss map

Recall that a minimal surface $S \subset \mathbb{R}^3$ is by definition a critical point of the area functional for compactly supported perturbations. Let $X: \Sigma \rightarrow \mathbb{R}^3$ be a conformal parametrization of S and $N: \Sigma \rightarrow S^2$ its Gauss map. If D is a compact subdomain of Σ , and $X_t: D \rightarrow \mathbb{R}^3$ is a normal variation of D with variation vector-field fN , $f \in C_0^\infty(D)$, the second derivative of area is given by

$$\frac{d^2}{dt^2} \text{Area}(X_t) = \int_D (|\nabla f|^2 + 2Kf^2) dA =: Q_D(f, f), \quad (7.1)$$

where K is the Gauss curvature. The second variation operator is defined to be

$$L =: -\Delta + 2K, \quad (7.2)$$

in terms of which

$$Q_D(f, f) = \int_D fLf dA.$$

The sign of Δ is determined by our definition in (2.3). The index of L on D , denoted $\text{ind}(D)$, is defined to be the dimension of the maximal subspace of $C_0^\infty(D)$ on which Q_D is negative definite. A domain $X(D) \subset S$ is said to be *stable* if $\text{ind}(D) = 0$, i.e., if (7.1) is nonnegative for all $f \in C_0^\infty(D)$.

For a non-compact minimal surface $X(\Sigma) = S \subset \mathbb{R}^3$, the index (Σ) is defined to be the supremum of $\text{ind}(D)$ taken over all compact $D \subset \Sigma$. The following theorem summarizes the results of several authors relating index to stability.

Theorem 7.1 *Let $X: \Sigma \rightarrow \mathbb{R}^3$ be a conformal minimal immersion of an orientable surface.*

- i) (Schwarz [67].) If the Gaussian image of Σ lies in a hemisphere, S is stable;*
- ii) (Barbosa and do Carmo [1].) If the area of the Gaussian image satisfies $A(N(\Sigma)) < 2\pi$, then S is stable. (Here $A(N(\Sigma))$ is the area of the Gaussian image disregarding multiplicities);*
- iii) (Fischer-Colbrie and Schoen [23], do Carmo and Peng [18], Pogorelov [61].) If S is complete, S is stable if and only if S is a plane;*
- iv) (Fischer-Colbrie [22], Gulliver [25], Gulliver and Lawson [26].) $\text{Index}(\Sigma) < \infty$ if and only if $\int_\Sigma |K| dA < \infty$.*

Without going too much into detail, we will give some of the background to the last two parts of the theorem above. The Gauss curvature K can be expressed as $-|\nabla N|^2/2$ and therefore we may write $L = -(\Delta + |\nabla N|^2)$. Observe that $Q_D(f, f) = \int_D (|\nabla f|^2 - |\nabla N|^2 f^2) dA$ is invariant under conformal change of metric and therefore $\text{ind}(D)$ depends only on N .

We may therefore define, for a meromorphic mapping $\Phi: \Sigma \rightarrow S^2$, the index $ind_\Phi(D)$ of a compact subdomain $D \subset \Sigma$ to be the number of negative eigenvalues of $L_\Phi = -(\Delta + |\nabla \Phi|^2)$, with Dirichlet boundary conditions on D . By definition, $ind(D) = ind_N(D)$. In particular if $\bar{\Sigma}$ is a compact Riemann surface and Φ a meromorphic function on $\bar{\Sigma}$,

$$index(\Phi) := ind_\Phi(\bar{\Sigma}) \quad (7.3)$$

is an integer invariant of $\Phi: \bar{\Sigma} \rightarrow S^2$.

For a complete immersed minimal surface $S \subset \mathbb{R}^3$, the condition of finite total curvature is equivalent to the conditions that the underlying Riemann surface Σ is a compact Riemann surface $\bar{\Sigma}$ with a finite number of points removed, and that the Gauss map N extends to a meromorphic function on $\bar{\Sigma}$ (See Theorem 2.2). Thus for any compact subdomain $D \subset \Sigma$,

$$ind(D) = ind_N(D) \leq index(N),$$

In particular the condition of finite total curvature implies that

$$index(\Sigma) = \sup\{ind(D) \mid D \subset \Sigma, D \text{ compact}\}$$

is finite. This is the easier implication of part *iii*) of Theorem 7.1. In fact these two indices are equal:

Proposition 7.1 ([22]) *Let $X: \Sigma \rightarrow \mathbb{R}^3$ be a complete immersed minimal surface of finite total curvature. Let $N: \Sigma \rightarrow S^2$ be its Gauss map. Then $ind(\Sigma) = index(N)$.*

As a consequence of this Proposition, statement *iii*) of Theorem 7.1 follows from the fact that the index of any nonconstant meromorphic function $\Phi: \bar{\Sigma} \rightarrow S^2$ is nonzero.

Remark 7.1 *Let ds_N^2 be the metric produced by pulling back the standard metric on S^2 by the Gauss map $N: \Sigma \rightarrow S^2$. If ds^2 denotes the metric induced by $X: \Sigma \rightarrow \mathbb{R}^3$, we have $ds_N^2 = \frac{1}{2}|\nabla N|^2 ds^2 = -K ds^2$. This metric is regular, away from the isolated zeros of K , and has constant curvature 1. The volume of $\bar{\Sigma}$ in this metric is therefore $4\pi \text{degree}(\Phi)$. Because the Dirichlet integrand $|\nabla f|^2 dA$ is invariant under conformal change of metric*

$$Q(f, f) = \int (|\nabla f|^2 + 2K f^2) dA = \int (|\tilde{\nabla} f|^2 - 2f^2) d\tilde{A},$$

where $\tilde{\nabla} f$ and $d\tilde{A}$ are computed in the metric ds_N^2 .

The associated differential operator is $\tilde{L} = -(\tilde{\Delta} + 2)$. Eigenfunctions of $\tilde{\Delta}$ can be defined by the standard variational procedure [54, 71]. The spectrum of $\tilde{\Delta}$ is discrete, infinite and nonnegative and we may think of $index(\Phi)$ as the number of eigenvalues of $\tilde{\Delta}$ that are strictly less than two.

It is natural to try to compute the index of stability of the minimal surfaces mentioned in this survey. We have seen that this is equivalent to computing the index of their extended Gauss maps. For the simplest case of the identity map from S^2 to itself, the index is just the number of eigenvalues of $\tilde{\Delta} = \Delta$ (the spherical Laplacian) less than two. Since the first nonzero eigenvalue is equal to 2 and $\lambda_0 = 0$ is a simple eigenvalue, it follows from Remark 7.1 above that $\text{index}(Id) = 1$.

Proposition 7.2 *The index of stability of Enneper's surface and the catenoid is equal to one.*

This follows immediately from the fact that the extended Gauss map of these genus-zero surfaces is a conformal diffeomorphism to the sphere. Osserman [58, 59] proved that the catenoid and Enneper's surface were the only complete minimal surfaces with this property. It is natural to ask whether or not index one characterizes these surfaces among all complete minimal surfaces of finite total curvature. The following result of Montiel-Ros ([54]) can be used to resolve this question.

Theorem 7.2 *Suppose $\Phi: \bar{\Sigma} \rightarrow S^2$ is a nonconstant holomorphic map of degree d defined on a compact Riemann surface of genus k .*

- i) If $k = 0$, $\text{index}(\Phi) = 1 \iff d = 1$.*
- ii) If $k = 1$, $\text{index}(\Phi) \geq 2$.*
- iii) If $k \geq 2$ then, for a generic conformal structure, $\text{index}(\Phi) \geq 2$.*
- iv) If $\text{index}(\Phi) = 1$, $d \leq 1 + [\frac{1+k}{2}]$.*

Statements *i)* and *ii)* are straightforward. Statements *iii)* and *iv)* follow from more general results in [54] (see Corollary 8 and Theorem 6) and the Brill-Noether Theorem.

Now suppose that S is a complete minimal surface whose index of stability is equal to 1. From (2.9), we have for the Gauss map N of a complete minimal surface $X: \Sigma \rightarrow \mathbb{P}^3$ of genus k

$$d = \text{degree}(N) \geq k + 1.$$

Combined with statement *iv)* of Theorem 7.2, this implies $k \leq 1$. Then statements *i)* and *ii)* imply that $k = 0$ and $d = 1$. As noted above, the catenoid and Enneper's surface are the only possibilities. Thus we have

Corollary 7.1 (Corollary 9 of [54] [50]) *A complete minimal surface with stability index equal to one must be the catenoid or Enneper's surface.*

Remark 7.2 *Montiel and Ros conjectured that it was possible to have a hyperelliptic Riemann surface whose two-fold covering map to S^2 had index equal to one, provided the branch values were sufficiently well distributed. This was proved by Souam [68].*

Theorem 7.3 *Given $\varepsilon > 0$, choose $\mathcal{P} = \{p_1 \dots p_{n_\varepsilon}\} \subset S^2$ such that the open ε -disks about the p_i are disjoint, but maximal in the sense that their complement does not contain an ε -disk. Suppose Σ is a hyperelliptic Riemann surface and $\phi: \Sigma \rightarrow S^2$ a two-fold covering of S^2 whose branch values are precisely \mathcal{P} . Then if $\varepsilon > 0$ is small enough, $\text{index}(\Phi) = 1$.*

The following results allows us to compute the index of stability of the Meeks-Jorge n -noids, the Chen-Gackstatter surface and its generalizations to higher symmetry (See Remark 4.2).

Proposition 7.3 (Montiel and Ros [54]) *Suppose $\phi: \overline{\Sigma} \rightarrow S^2$ is a degree $d \geq 1$ holomorphic map. If all the branch value of ϕ lie on a great circle*

$$\text{index}(\Phi) = 2d - 1. \quad (7.4)$$

Choe [12], proved a weaker version of this proposition.

Theorem 7.4 ([54]) *The index of stability of:*

- i) the n -noid is $2n - 3$;*
- ii) the Chen-Gackstatter surface is 3.*

Proof. From Section 2.2 we know that the Gauss map of the Chen-Gackstatter surface has degree $d = 2$, and all of its branch values are real. The n -noid $n \geq 2$ can be produced with the Weierstrass data $\overline{\Sigma} = S^2$,

$$g = z^{n-1}$$

$$dh = (z^n + z^{-n} + 2)z^{-1}dz.$$

(When $n = 2$ we have the catenoid.) In both cases we can apply Proposition 7.3. \square

The equivalence of finite total curvature and finite index suggests that there should be an explicit relationship between d , the degree of the Gauss map, and the index of stability. Tysk [71] was the first to show that the index could be estimated by the degree:

$$\frac{\text{Index}(N)}{\text{degree}(N)} < c \text{ where } c \sim 7.68.$$

Choe [12] conjectured that a weakened version of (7.3) was true in general for all complete minimal surfaces M of finite total curvature $-4\pi d$:

Conjecture 7.1

$$\text{index}(M) \leq 2d - 1 \quad (7.5)$$

The conjectured inequality (7.5) cannot be replaced in general by an equality. Nayatani (private communication) has pointed out to us that an example written down by Kusner [44], Rosenberg-Toubiana [65], R. Bryant [6, 7] with 4 flat ends, genus 0, and Gauss map of degree $d = 3$, gives strict inequality. On any surface the support function $u = X \cdot N$ satisfies $Lu = 0$, where L is defined in (7.2), and because the ends are flat, u is bounded. Using this fact, it can be shown that the index of this surface is 4. Since $d = 3$, $\text{index} = 4 < 5 = 2d - 1$.

Nayatani [55] computed the index of the Costa surface, M_2 of Theorem 3.2. This was done by computing the index of its Gauss map, which is the derivative of the Weierstrass \wp -function, geometrically normalized, on the square torus. The geometric normalization does not change the index.

He subsequently improved that result:

Theorem 7.5 (Nayatani [56]) *Let M_k be the Hoffman-Meeks surface of genus $k - 1$, described in Theorem 3.2. For $2 \leq k \leq 38$,*

$$\text{index}(M_k) = 2k + 1.$$

The Gauss map of M_k has degree $k + 1$, so equality holds in (7.5) for these surfaces. For genus 0 surfaces Montiel and Ros [54] and Ejiri and Kotani [20] independently proved the following result.

Theorem 7.6 *Formula (7.4) holds for a generic complete, genus zero, minimal surface of total curvature $-4\pi d$. In general, $\text{index}(M) \leq 2d - 1$ for such a surface.*

This result holds even for branched minimal surfaces and involves an analysis of bounded Jacobi fields.

We end this chapter with a discussion of the work of Choe [12] about the *vision number* of a minimal surface with respect to a vector field V on \mathbb{R}^3 . For a vector field V on $S \subset \mathbb{R}^3$, Choe defines the *horizon* of V to be

$$H(S, V) = \{s \in S \mid V \in T_s S\},$$

and the vision number, to be

$$v(S, V) = \# \text{ components of } S - H(S, V).$$

Suppose V is a variation vector field associated with a family of parallel translations, or rotations about a line. (Both are Killing vector fields on \mathbb{R}^3). Consider the restriction of such a vector field to S , and let \widehat{V} be its projection onto the normal bundle of S . Then \widehat{V} is a Jacobi field. That is, $\widehat{V} \cdot N$ is in the kernel of $L = -\Delta + 2K$. The same holds for the position vector field $V_0(p) = \overline{p - p_0}$, for any fixed origin $p_0 \in \mathbb{R}^3$. This can be used to show that any connected subset of $S - H(S, V)$ is stable, for any one of these vector fields V . Choe uses this to prove

Theorem 7.7 *Let $S \subset \mathbb{R}^3$ be a complete minimal surface of finite total curvature $-4\pi d$. Let V_T, V_R denote the Killing vector fields generated by translation and rotation, respectively. Then*

$$i) \text{ Index}(S) \geq v(S, V_T) - 1;$$

$$ii) \text{ } d \geq v(S, V_T)/2;$$

If all the ends are parallel

$$iii) \text{ Index}(S) \geq v(S, V_R) - 1;$$

If all the ends are embedded and parallel

$$iv) \text{ } d \geq v(S, V_R)/2.$$

From statement *iii)* of the above theorem we can get a lower bound for the index of the surfaces M_k of Theorem 3.2. Observe that the same is true for the $M_{k,\alpha}$ of Section 4. These surfaces have genus $k - 1$, parallel embedded ends and k vertical planes of reflective symmetry, which intersect in a vertical line, \mathcal{L} , orthogonal to the ends, and which divide the surface into $2k$ regions. Their Gauss maps have degree $d = k + 1$. If V_R is the Killing field generated by rotation about \mathcal{L} , then it is clear that $H(M_{k,\alpha}, V_R)$ contains the intersection of $M_{k,\alpha}$ with each plane of symmetry. Hence $v(M_{k,\alpha}, V_R) \geq 2k$. By Theorem 7.7 *ii)* we have

Corollary 7.2 *The index of stability of the surfaces $M_k = M_{k,0}$ and $M_{k,\alpha}$ satisfies*

$$\text{Index}(M_{k,\alpha}) \geq 2k - 1 = 2(\text{genus } M_{k,\alpha}) + 1. \quad (7.6)$$

Remark 7.3 *The symmetry of the surfaces $M_{k,\alpha}$ imply that $v(M_{k,\alpha}, V_R)$ is an integer multiple of $2k$. But by statement *iv)* of Theorem 7.7, $2(k + 1) = 2d \geq v(M_{k,\alpha}, V_R)$. Hence $v(M_{k,\alpha}, V_R) = 2k$. Thus this particular argument cannot be improved to give a better lower bound. As we know from Theorem 7.5, $\text{index}(M_k) = 2k + 1, 2 \leq k \leq 38$, so this estimate is not sharp in this range.*

References

- [1] J. L. Barbosa and M. do Carmo. On the size of a stable minimal surface in \mathbb{R}^3 . *American Journal of Mathematics*, 98(2):515–528, 1976.
- [2] L. Barbosa and A. G. Colares. *Minimal Surfaces in \mathbb{R}^3* . Springer-Verlag, 1986. Lecture notes in mathematics, No. 1195.
- [3] D. Bloß. *Elliptische Funktionen und vollständige Minimalflächen*. PhD thesis, University of Berlin, 1989.
- [4] E. Boix and L. Capdequi. Deformation families of complete minimal surfaces with four catenoid ends. (Video tape) GANG, August 1991.
- [5] E. Boix and M. Wohlgemuth. Numerical results on embedded minimal surfaces of finite total curvature. In preparation.
- [6] R. Bryant. A duality theorem for Willmore surfaces. *Journal of Differential Geometry*, 20:23–53, 1984.
- [7] R. Bryant. Surfaces in conformal geometry. In *The Mathematical Heritage of Hermann Weyl*, volume 48 of *Proceedings of Symposia in Pure Math.* AMS, 1988. R. O. Wells, editor.
- [8] M. Callahan, D. Hoffman, and J. Hoffman. Computer graphics tools for the study of minimal surfaces. *Communications of the ACM*, 31(6):648–661, 1988.
- [9] M. Callahan, D. Hoffman, and W. H. Meeks III. Embedded minimal surfaces with an infinite number of ends. *Inventiones Math.*, 96:459–505, 1989.
- [10] C. C. Chen and F. Gackstatter. Elliptic and hyperelliptic functions and complete minimal surfaces with handles. *Instituto de Matemática e Estatística-Universidade de São Paulo*, 27, 1981.
- [11] C. C. Chen and F. Gackstatter. Elliptische und Hyperelliptische Functionen und vollständige Minimalflächen von Enneperschen Typ. *Math. Ann.*, 259:359–369, 1982.
- [12] J. Choe. Index, vision number, and stability of complete minimal surfaces. *Arch. Rational Mech. Anal.*, 109(3):195–212, 1990.
- [13] T. Choi, W. H. Meeks III, and B. White. A rigidity theorem for properly embedded minimal surfaces in \mathbb{R}^3 . *Journal of Differential Geometry*. March, 1990.
- [14] C. Costa. *Imersões mínimas em \mathbb{R}^3 de gênero um e curvatura total finita*. PhD thesis, IMPA, Rio de Janeiro, Brasil, 1982.

- [15] C. Costa. Example of a complete minimal immersion in \mathbb{R}^3 of genus one and three embedded ends. *Bull. Soc. Bras. Mat.*, 15:47–54, 1984.
- [16] C. Costa. Uniqueness of minimal surfaces embedded in \mathbb{R}^3 with total curvature 12π . *Journal of Differential Geometry*, 30(3):597–618, 1989.
- [17] U. Dierkes, S. Hildebrandt, A. Küster, and O. Wohlrab. *Minimal Surfaces I*. Grundlehren der mathematischen Wissenschaften 296. Springer-Verlag, 1992.
- [18] M. do Carmo and C. K. Peng. Stable minimal surfaces in \mathbb{R}^3 are planes. *Bulletin of the AMS*, 1:903–906, 1979.
- [19] N. do Espírito-Santo. *Superfícies mínimas completas em \mathbb{R}^3 com fim de tipo Enneper*. PhD thesis, University of Niteroi, Brazil, 1992.
- [20] N. Ejiri and M. Kotani. Minimal surfaces in $S^{2m}(1)$ with extra eigenfunctions. *Quart. J. Math. Oxford, Second Series*, 172:421–440, 1992.
- [21] Y. Fang and W. H. Meeks III. Some global properties of complete minimal surfaces of finite topology in \mathbb{R}^3 . *Topology*, 30(1):9–20, 1991.
- [22] D. Fischer-Colbrie. On complete minimal surfaces with finite Morse index in 3-manifolds. *Inventiones Math.*, 82:121–132, 1985.
- [23] D. Fischer-Colbrie and R. Schoen. The structure of complete stable minimal surfaces in 3-manifolds of non-negative scalar curvature. *Comm. on Pure and Appl. Math.*, 33:199–211, 1980.
- [24] F. Gackstatter. Über die Dimension einer Minimalfläche und zur Ungleichung von St. Cohn-Vossen. *Arch. Rational Mech. Anal.*, 61(2):141–152, 1976.
- [25] R. Gulliver. Index and total curvature of complete minimal surfaces. *Proc. Symp. Pure Math.*, 44:207–212, 1986.
- [26] R. Gulliver and H. B. Lawson. The structure of minimal hypersurfaces near a singularity. *Proc. Symp. Pure Math.*, 44:213–237, 1986.
- [27] J.-J. Hadamard. Les surfaces à courbures opposées et leurs lignes géodésiques. *Journal de Mathématiques Pures et Appliquées*, 4:27–73, 1898.
- [28] D. Hoffman. The discovery of new embedded minimal surfaces: Elliptic functions, symmetry; computer graphics. In *Proceedings of the Berlin Conference on Global Differential Geometry*, Springer Lecture Series 1748, 1985.
- [29] D. Hoffman. The computer-aided discovery of new embedded minimal surfaces. *Mathematical Intelligencer*, 9(3):8–21, 1987.

- [30] D. Hoffman, H. Karcher, and W. H. Meeks III. One-parameter families of embedded complete minimal surfaces of finite topology. *GANG preprint* in preparation.
- [31] D. Hoffman, H. Karcher, and H. Rosenberg. Embedded minimal annuli in \mathbb{R}^3 bounded by a pair of straight lines. *Comment. Math. Helvetici*, 66:599–617, 1991.
- [32] D. Hoffman, H. Karcher, and F. Wei. Adding handles to the helicoid. *Bulletin of the AMS, New Series*, 29(1):77–84, 1993.
- [33] D. Hoffman, H. Karcher, and F. Wei. The genus one helicoid and the minimal surfaces that led to its discovery. In *Global Analysis and Modern Mathematics*. Publish or Perish Press, 1993. K. Uhlenbeck, editor, p. 119–170.
- [34] D. Hoffman and W. H. Meeks III. A variational approach to the existence of complete embedded minimal surfaces. *Duke Journal of Math.*, 57(3):877–894, 1988.
- [35] D. Hoffman and W. H. Meeks III. The asymptotic behavior of properly embedded minimal surfaces of finite topology. *Journal of the AMS*, 2(4):667–681, 1989.
- [36] D. Hoffman and W. H. Meeks III. Embedded minimal surfaces of finite topology. *Annals of Mathematics*, 131:1–34, 1990.
- [37] D. Hoffman and W. H. Meeks III. Les surfaces minimales: la caténoïde par les deux bouts. *Quadrature*, 5:31–47, 1990.
- [38] D. Hoffman and W. H. Meeks III. Limits of minimal surfaces and Scherk’s Fifth Surface. *Arch. Rat. Mech. Anal.*, 111(2):181–195, 1990.
- [39] D. Hoffman and W. H. Meeks III. Minimal surfaces based on the catenoid. *Amer. Math. Monthly, Special Geometry Issue*, 97(8):702–730, 1990.
- [40] D. Hoffman and W. H. Meeks III. The strong halfspace theorem for minimal surfaces. *Inventiones Math.*, 101:373–377, 1990.
- [41] J. T. Hoffman. MESH manual. *GANG preprint series II*, #35.
- [42] L. Jorge and W. H. Meeks III. The topology of complete minimal surfaces of finite total Gaussian curvature. *Topology*, 22(2):203–221, 1983.
- [43] H. Karcher. Construction of minimal surfaces. *Surveys in Geometry*, pages 1–96, 1989. University of Tokyo, 1989, and Lecture Notes No. 12, SFB256, Bonn, 1989.
- [44] R. Kusner. Conformal geometry and complete minimal surfaces. *Bulletin of the AMS*, 17(2):291–295, 1987.

- [45] R. Kusner. *Global geometry of extremal surfaces in three-space*. PhD thesis, University of California, Berkeley, 1988.
- [46] R. Kusner. Bubbles, conservation laws and balanced diagrams. In *Geometric Analysis and Computer Graphics*, volume 17 of *MSRI Publications*. Springer-Verlag, 1990. P. Concus and R. Finn and D. Hoffman, editors.
- [47] H. B. Lawson. Complete minimal surfaces in S^3 . *Annals of Math.*, 92:335–374, 1970.
- [48] H. B. Lawson, Jr. *Lectures on Minimal Submanifolds*. Publish or Perish Press, Berkeley, 1971.
- [49] F. J. Lopez. The classification of complete minimal surfaces with total curvature greater than -12π . *Trans. Amer. Math. Soc.*, 334(1):49–74, 1992.
- [50] F. J. Lopez and A. Ros. Complete minimal surfaces with index one and stable constant mean curvature surfaces. *Comm. Math. Helv.*, 64:34–43, 1989.
- [51] F. J. Lopez and A. Ros. On embedded complete minimal surfaces of genus zero. *Journal of Differential Geometry*, 33(1):293–300, 1991.
- [52] W. H. Meeks III and H. Rosenberg. The maximum principle at infinity for minimal surfaces in flat three-manifolds. *Commentari Mathematici Helvetici*, 65:255–270, 1990.
- [53] W. H. Meeks III and H. Rosenberg. The geometry and conformal structure of properly embedded minimal surfaces of finite topology in E^3 . *Inventiones*, 114:625–639, 1993.
- [54] S. Montiel and A. Ros. Schrödinger operators associated to a holomorphic map. In *Global Differential Geometry and Global Analysis (Berlin, 1990)*, volume 1481 of *Lecture Notes in Mathematics*, pages 147–174. Springer-Verlag, 1990.
- [55] S. Nayatani. On the Morse index of complete minimal surfaces in Euclidean space. *Osaka J. Math.*, 27:441–451, 1990.
- [56] S. Nayatani. Morse index of complete minimal surfaces. In *The Problem of Plateau*, pages 181–189. World Scientific Press, 1992. Th. M. Rossi, editor.
- [57] J. C. C. Nitsche. A characterization of the catenoid. *Journal of Math. Mech.*, 11:293–302, 1962.
- [58] R. Osserman. Global properties of minimal surfaces in E^3 and E^n . *Annals of Math.*, 80(2):340–364, 1964.
- [59] R. Osserman. *A Survey of Minimal Surfaces*. Dover Publications, New York, 2nd edition, 1986.

- [60] J. Pérez and A. Ros. Some uniqueness and nonexistence theorems for embedded minimal surfaces. *Math. Ann.*, 295(3):513–525, 1993.
- [61] A. V. Pogorelov. On the stability of minimal surfaces. *Soviet Math. Dokl.*, 24:274–276, 1981.
- [62] B. Riemann. Über die Fläche vom kleinsten Inhalt bei gegebener Begrenzung. *Abh. Königl. d. Wiss. Göttingen, Mathem. Cl.*, 13:3–52, 1867.
- [63] B. Riemann. *Oeuvres Mathématiques de Riemann*. Gauthiers-Villars, Paris, 1898.
- [64] H. Rosenberg and E. Toubiana. A cylindrical type complete minimal surface in a slab of \mathbb{R}^3 . *Bull. Sc. Math. III*, pages 241–245, 1987.
- [65] H. Rosenberg and E. Toubiana. Remarks on deformations of minimal surfaces. *Transactions of the AMS*, 1987.
- [66] R. Schoen. Uniqueness, symmetry, and embeddedness of minimal surfaces. *Journal of Differential Geometry*, 18:791–809, 1983.
- [67] H. A. Schwarz. *Gesammelte Mathematische Abhandlungen*, volume 1. Springer, Berlin, 1890.
- [68] R. Souam. Exemples d’applications holomorphes d’indice un. *Ann. Inst. Fourier (Grenoble)*, 2:369–381, 1993.
- [69] E. Thayer. Dissertation in preparation.
- [70] E. Thayer. Generalized Chen-Gackstatter minimal surfaces, Scherk’s singly-periodic minimal surface and Riemann surfaces of infinite genus. *GANG preprint*, 1994.
- [71] J. Tysk. Eigenvalue estimates with applications to minimal surfaces. *Pacific J. of Math.*, 128:361–366, 1987.
- [72] M. Wohlgemuth. Higher genus minimal surfaces by growing handles out of a catenoid. *Manuscripta Math.*, 70:397–428, 1991.
- [73] M. Wohlgemuth. *Vollständige Minimalflächen höheren Geschlechts und endlicher Totalkrümmung*. PhD thesis, University of Bonn, April 1993.
- [74] M. Wohlgemuth. Higher genus minimal surfaces of finite total curvature. preprint, 1994.